

Stability Problems in Symbolic Integration^{*}

Shaoshi Chen

KLMM, Academy of Mathematics and Systems Science
Chinese Academy of Sciences, Beijing 100190, China
schen@amss.ac.cn

ABSTRACT

This paper aims at initializing a dynamical aspect of symbolic integration by studying stability problems in differential fields. We present some basic properties of stable elementary functions and D-finite power series that enable us to characterize three special families of stable elementary functions including rational functions, logarithmic functions, and exponential functions. Some problems for future studies are proposed towards deeper dynamical studies in differential and difference algebra.

CCS CONCEPTS

• Computing methodologies → Algebraic algorithms.

KEYWORDS

Dynamical systems, stable functions, symbolic integration, symbolic summation

ACM Reference Format:

Shaoshi Chen. 2022. Stability Problems in Symbolic Integration. In *Proceedings of the International Symposium on Symbolic and Algebraic Computation (ISSAC '22)*, July 4–7, 2022, Lille, France. ACM, New York, NY, USA, 8 pages. <https://doi.org/>

1 INTRODUCTION

In the proofs of the irrationality and transcendence of e and π , various kinds of definite integrals such as

$$\int_{-1}^1 (1-x^2)^n \cos(xz) dx \quad \text{and} \quad \int_0^\pi \frac{x^n (a-bx)^n}{n!} \sin(x) dx$$

are used [32, 33]. In the process of deriving linear recurrences in n for these integrals, one may realize that the integrands are so nice that their shape is stable under indefinite integration, which means the integrands can be integrated arbitrarily often in terms of elementary functions. Some typical functions with this nice property are polynomials, radicals of the form $(x-c)^r$ with $c \in \mathbb{C}$ and $r \in \mathbb{Q}$ and basic transcendental elementary functions such as $\log(x)$, $\exp(x)$, $\sin(x)$, $\cos(x)$, etc.. One may be curious about whether these are the only possible functions that have this feature. This motivates our dynamical thinking in symbolic integration.

^{*}S. Chen was partially supported by the NSFC grants 11871067, 12288201, and the Fund of the Youth Innovation Promotion Association, CAS.

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for components of this work owned by others than the author(s) must be honored. Abstracting with credit is permitted. To copy otherwise, or republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee. Request permissions from permissions@acm.org.

ISSAC '22, July 4–7, 2022, Lille, France

© 2022 Copyright held by the owner/author(s). Publication rights licensed to ACM.

ACM ISBN ... \$15.00

<https://doi.org/>

The problem of integration in finite terms or in closed form is one of the oldest problems in calculus. Since the initial work of Liouville from 1832 to 1841, different systematical approaches have been developed to study this problem [19, 23, 35, 40]. A comprehensive historical study of Liouville's work on integration in finite terms is given in [26] and also some gentle introductory notes are [20, 27]. After the birth of differential algebra, the problem is formulated in a pure algebraic fashion and then solved by Risch [38, 39] with further developments in symbolic integration [7, 8, 11, 14, 31, 36, 42, 46, 49]. The aim of symbolic integration is developing practical algorithms and software for solving the integration problem of elementary functions and other more general special functions from mathematical physics. The standard references on symbolic integration are firstly Bronstein's book [8] and some chapters in [12, 50]. Recently, Raab gave an informative survey [37] on the Risch algorithm and its recent developments. Another significant current trend is the arithmetic studies of elementary integration initialized by Masser and Zannier in [10, 29, 30, 51].

Self-maps on structured sets are ubiquitous in mathematics. The theory of dynamical systems is the mathematics of self-maps. So the dynamical way of thinking has inspired many interdisciplinary areas in mathematics such as arithmetic dynamics in number theory [45] and complex dynamics in complex analysis and geometry [5]. The fundamental objects in differential algebra are differential fields and their extensions. The indefinite integration problem of elementary functions can be formulated in terms of differential fields. In this paper, we first view differential fields as dynamical systems where derivations play the role of self-maps and then investigate the dynamical aspect of elementary integration by determining which elementary functions can be iteratively integrated in terms of elementary functions infinite times (see the precise formulation in Problem 3.5). Through the dynamical lens, we will see some new landscape of symbolic integration and also understand better why the integrals from the beginning are so nice.

The remainder of this paper is organized as follows. We recall some basic terminologies in dynamical systems and differential algebra and then define stability problems in Section 2. After bridging the connection, we explore the basic properties of stable elementary functions and characterize three families of special stable elementary functions in Section 3. In Section 4, we study the stability problem on D-finite powers series and then conclude our paper in 5 by proposing some problems for future research.

2 STABILITY IN DIFFERENTIAL FIELDS

We first bridge the connection between dynamical systems and differential algebra. A (discrete) *dynamical system* is a pair (A, ϕ) with A being a set and $\phi : A \rightarrow A$ being a self-map on A . We recall the definition of four special subsets that are crucial for understanding a dynamical system.

DEFINITION 2.1. Let (A, ϕ) be a dynamical system and $a \in A$. Then we say that

- (1) the element a is a fixed point of ϕ if $\phi(a) = a$. The set of all fixed points is denoted by $\text{Fix}(\phi, A)$;
- (2) the element a is a periodic point of ϕ if $\phi^n(a) = a$ for some positive $n \in \mathbb{N}$. The set of all periodic points is denoted by $\text{Per}(\phi, A)$;
- (3) the element a is stable in the system (A, ϕ) if there exists a sequence $\{a_i\}_{i \geq 0}$ in A such that $a_0 = a$ and $\phi(a_{i+1}) = a_i$ for all $i \in \mathbb{N}$. The set of all stable elements is denoted by $\text{Stab}(\phi, A)$;
- (4) the element a is attractive in the system (A, ϕ) if for any $i \in \mathbb{N}$, there exists $a_i \in A$ such that $a = \phi^i(a_i)$. The set of all attractive elements is denoted by $\text{Attrac}(\phi, A)$ that is equal to $\bigcap_{i \in \mathbb{N}} \phi^i(A)$.

For any dynamical system (A, ϕ) , we have the inclusions:

$$\text{Fix}(\phi, A) \subseteq \text{Per}(\phi, A) \subseteq \text{Stab}(\phi, A) \subseteq \text{Attrac}(\phi, A).$$

It is not hard to see that the first three inclusions may be proper by definition. For the last one, Godelle in [15] presented a concrete example as below and also discussed the question of deciding whether the inclusion $\text{Stab}(\phi, A) \subseteq \text{Attrac}(\phi, A)$ is indeed an equality in various settings.

EXAMPLE 2.2 (GODELLE'S EXAMPLE). Let $A = \{(i, j) \in \mathbb{Z}^2 \mid 0 \leq j \leq \max(i-1, 0)\}$ and let $\phi : A \rightarrow A$ be defined by $\phi(i, j) = (i, j-1)$ for positive j , and $\phi(i, 0) = (\min(i-1, 0), 0)$. Then the subset $\text{Stab}(\phi, A)$ is empty, but the subset $\text{Attrac}(\phi, A)$ is $\{(i, 0) \mid i \leq 0\}$.

For the convenience of later usage, we summarize the results on stable and attractive subsets in [15, Sections 2-3] as follows.

THEOREM 2.3 (GODELLE'S THEOREM). Let A be a set and $\phi : A \rightarrow A$ be a self-map on A . Then

- (i) The stable subset $\text{Stab}(\phi, A)$ is the largest one among all subsets of A satisfying the property $\phi(B) = B$ for $B \subseteq A$.
- (ii) If ϕ is either surjective on A or injective on $\phi^n(A)$ for some $n \in \mathbb{N}$, then $\text{Stab}(\phi, A) = \text{Attrac}(\phi, A)$.
- (iii) If A is an infinite-dimensional vector space over a field k , and ϕ is a linear map such that $\dim_k(\text{Ker}(\phi)) = 1$. Then $\text{Stab}(\phi, A) = \text{Attrac}(\phi, A)$.

Throughout this paper, we assume that all fields are of characteristic zero. For any field K , we call an additive map $\delta : K \rightarrow K$ a derivation on K if $\delta(fg) = g\delta(f) + f\delta(g)$ for all $f, g \in K$. By induction, we have the general Leibniz formula

$$\delta^n(fg) = \sum_{i=0}^n \binom{n}{i} \delta^i(f) \delta^{n-i}(g) \quad (2.1)$$

for all $n \in \mathbb{N}$ and $f, g \in K$. We call the pair (K, δ) a differential field. An element $c \in K$ is called a constant in K if $\delta(c) = 0$. The set of all constants in K forms a subfield of K , denoted by C_K . Note that δ is a C_K -linear map on K . An element $f \in K$ is said to be integrable in K if there exists $g \in K$ such that $f = \delta(g)$. By the C_K -linearity of the derivation δ , we see that the set $\delta(K)$ of all integrable elements in K forms a linear subspace of K over C_K . A differential field (K_1, δ_1) is called a differential extension of the differential field (K, δ) if $K \subseteq K_1$ and $\delta_1|_K = \delta$. By abuse of notation, we use the same symbol for derivations on differential extensions.

The ring of linear differential operators over the differential field (K, δ) is denoted by $K\langle D \rangle$, in which we have the commutation rule

$$D \cdot f = f \cdot D + \delta(f) \quad \text{for all } f \in K.$$

One of basic properties of the ring $K\langle D \rangle$ is that it is a left Euclidean domain, in which we can define and effectively compute the greatest common right divisor (GCRD) and least common left multiple (LCLM) of polynomials if K is a computable field. For more results on this ring, one can see [1, 9, 34]. The general Leibniz rule (2.1) now is translated into the general commutation rule in $K\langle D \rangle$ as follows

$$D^n \cdot f = \sum_{i=0}^n \binom{n}{i} \delta^i(f) \cdot D^{n-i} \quad (2.2)$$

for all $n \in \mathbb{N}$ and $f \in K$. Similarly, we have the following formula

$$f \cdot D^n = \sum_{i=0}^n (-1)^i \binom{n}{i} D^{n-i} \cdot \delta^i(f) \quad (2.3)$$

for all $n \in \mathbb{N}$ and $f \in K$. The differential field K can be viewed as a $K\langle D \rangle$ -module via the action

$$L(f) = \sum_{i=0}^d \ell_i \delta^i(f)$$

for any $L = \sum_{i=0}^d \ell_i D^i \in K\langle D \rangle$ and $f \in K$.

We now view the differential field (K, δ) as a dynamical system and consider its stable and attractive subsets. By the C_K -linearity of the derivation δ , it is easy to show that the sets of all stable and attractive elements in K form linear subspaces of K over C_K , respectively.

PROPOSITION 2.4. Let (K, δ) be a differential field. Then

$$\text{Stab}(\delta, K) = \text{Attrac}(\delta, K).$$

PROOF. Let C_K be the constant subfield of K . Then K can be viewed as a vector space over C_K . If there exists an $f \in K$ such that $\delta(f) \neq 0$, then f must be transcendental over C_K by Lemma 3.3.2 in [8, p. 86]. So the dimension $\dim_{C_K}(K)$ is either one or infinite. When $\dim_{C_K}(K)$ is one, we have $K = C_K$ and then both $\text{Stab}(\delta, K)$ and $\text{Attrac}(\delta, K)$ are equal to $\{0\}$. So the equality holds. When $\dim_{C_K}(K)$ is infinite, we get the equality by Theorem 2.3 (iii), since δ is a linear map on K over C_K and $\dim_{C_K}(\text{Ker}(\delta)) = 1$. ■

So it is not needed to distinguish the stable elements and attractive elements in a dynamical system arising from a differential field. The central problem considered in this paper is as follows.

PROBLEM 2.5 (STABILITY PROBLEM). Given an element f in a differential field (K, δ) , decide whether f is stable in (K, δ) or not. More generally, describe the structure of the stable subset $\text{Stab}(\delta, K)$.

DEFINITION 2.6. A differential field (K, δ) is said to be regular if there exists an $x \in K$ such that $\delta(x) = 1$.

In a regular differential field, we have the usual formula

$$\delta^m(x^n) = n(n-1) \cdots (n-m+1)x^{n-m}$$

for all $m, n \in \mathbb{N}$ with $m \leq n$.

EXAMPLE 2.7. Let K be the field $\mathbb{C}(x)$ of rational functions in x over the field of complex numbers \mathbb{C} . If δ is the usual derivation d/dx , then (K, δ) is regular since $\delta(x) = 1$. If δ is the Eulerian derivation $x \cdot d/dx$, then (K, δ) is not regular since $1/x$ has no anti-derivative in $\mathbb{C}(x)$.

PROPOSITION 2.8. A differential field (K, δ) is regular if and only if all constants are stable in (K, δ) . Moreover, an element f in a regular differential field (K, δ) is stable if and only if $\delta(f)$ is stable.

PROOF. Suppose that (K, δ) is regular. Then there exists an $x \in K$ such that $\delta(x) = 1$. Let c be any constant in K . Note that

$$c = \delta^i \left(\frac{c}{i!} x^i \right) \quad \text{for any } i \in \mathbb{N}.$$

So $c \in \text{Attrac}(\delta, K) = \text{Stab}(\delta, K)$. The necessity is obvious since 1 is always a constant.

For the second assertion, suppose that f is stable in K . Then Theorem 2.3 (i) says that the stability is preserved under the derivation. So $\delta(f)$ is stable. Conversely, suppose that $\delta(f)$ is stable. Then there exists a sequence $\{g_i\}_{i \in \mathbb{N}}$ in K such that $g_0 = \delta(f)$ and $g_i = \delta(g_{i+1})$ for all $i \in \mathbb{N}$. It is clear that all of the g_i 's are stable by definition. By the equality $\delta(f) = \delta(g_1)$ and the additive property of the derivation δ , it follows that $\delta(f - g_1) = 0$, thus $f - g_1 = c$ for some $c \in C_K$, which yields $f = g_1 + c$. Since K is regular, c is also stable in K by the first assertion. Thus, f is stable in K . ■

REMARK 2.9. We point out that the regularity assumption in the above proposition is needed. In the differential field $(\mathbb{C}(x), x \cdot d/dx)$ which is not regular shown in Example 2.7, any nonzero constant c is not stable but its derivative is always stable.

LEMMA 2.10. Let (K, δ) be a regular differential field with $\delta(x) = 1$ and $f \in K$. Then

- (i) Let n be a positive integer. Then $f = \delta^n(g)$ for some $g \in K$ if and only if for all i with $0 \leq i \leq n-1$, there exists an $h_i \in K$ such that $x^i f = \delta(h_i)$.
- (ii) f is stable in (K, δ) if and only if for all $i \in \mathbb{N}$, $x^i f = \delta(g_i)$ for some $g_i \in K$.

PROOF. For showing the sufficiency of the first assertion, we suppose that $f = \delta^n(g)$ for some $g \in K$. Then the formula (2.3) implies that for all i with $0 \leq i \leq n-1$, we have

$$x^i \cdot D^n = \sum_{j=0}^n (-1)^j \binom{n}{j} D^{n-j} \cdot \delta^j(x^i).$$

Since $\delta^j(x^i) = 0$ if $j > i$, we have $x^i \cdot D^n = D \cdot L_i$ with

$$L_i = \sum_{j=0}^i (-1)^j j! \binom{n}{j} \binom{i}{j} D^{n-j-1} \cdot x^{i-j}.$$

Then $x^i f = x^i \delta^n(g) = (x^i \cdot D^n)(g) = \delta(h_i)$ with $h_i = L_i(g) \in K$.

We prove the necessity by induction on n . The assertion holds obviously in the base case when $n = 1$. We now assume that the assertion holds for $n < m$. Suppose that there exist elements $h_0, \dots, h_{m-1} \in K$ such that

$$f = \delta(h_0), \quad \dots, \quad x^{m-1} f = \delta(h_{m-1}).$$

The goal is to show that $f = \delta^m(g)$ for some $g \in K$. By the induction hypothesis, there exists an element $u \in K$ such that $f = \delta^{m-1}(u)$

using the first $m-1$ equalities $x^j f = \delta(h_j)$ for j with $0 \leq j \leq m-2$. By the formula (2.3), we have

$$x^{m-1} \cdot D^{m-1} = D \cdot L + (m-1)!$$

for some $L \in K\langle D \rangle$. We claim that we can choose

$$g = \frac{1}{(m-1)!} (h_{m-1} - L(u)).$$

Since $f = \delta^{m-1}(u)$ and $x^{m-1} f = \delta(h_{m-1})$, we have

$$\begin{aligned} D^{m-1} \cdot (x^{m-1} f) &= (D^{m-1} \cdot (x^{m-1} \cdot D^{m-1}))(u) \\ &= (D^{m-1} \cdot (DL + (m-1)!))(u) \\ &= D^m(L(u)) + (m-1)! D^{m-1}(u) \\ &= \delta^m(L(u)) + (m-1)! f. \end{aligned}$$

Since $D^{m-1} \cdot (x^{m-1} f) = \delta^{m-1}(\delta(h_{m-1})) = \delta^m(h_{m-1})$, we get $\delta^m(L(u)) + (m-1)! f = \delta^m(h_{m-1})$, which implies the claim.

The second assertion follows immediately from the first one. ■

THEOREM 2.11. Let (K, δ) be a regular differential field with $\delta(x) = 1$. Then the stable subset $\text{Stab}(\delta, K)$ forms a $C_K[x]$ -module and it is closed under differentiation.

PROOF. Since the derivation δ is a linear map on K over C_K , we have that $\delta^i(K)$ is a linear subspace of K for any $i \in \mathbb{N}$. By Proposition 2.4, $\text{Stab}(\delta, K) = \bigcap_{i \in \mathbb{N}} \delta^i(K)$ is also a linear subspace of K over C_K . To show that $\text{Stab}(\delta, K)$ forms a $C_K[x]$ -module, it suffices to prove that for any $f \in \text{Stab}(\delta, K)$ and $n \in \mathbb{N}$, we have $x^n f \in \text{Stab}(\delta, K)$. By Lemma 2.10, an element $g \in K$ is stable if and only if for all $i \in \mathbb{N}$, $x^i g = \delta(h_i)$ for some $h_i \in K$. Since f is stable, we have for all $j \in \mathbb{N}$, $x^j f = \delta(u_j)$ for some $u_j \in K$. This implies that for all $i, n \in \mathbb{N}$, $x^i(x^n f) = \delta(v_{i,n})$ for some $v_{i,n} \in K$. So $x^n f$ is stable for any $n \in \mathbb{N}$. The closure property of $\text{Stab}(\delta, K)$ under differentiation follows from Theorem 2.3 (i). ■

We now recall the notion of orders of rational functions and their basic properties from [8, Chapter 4]. Let F be a field of characteristic zero and $F(t)$ be the field of rational functions in t over F . Let $p \in F[t]$ be an irreducible polynomial and $f \in F(t)$. We can always write $f = p^m a/b$, where $m \in \mathbb{Z}$, $a, b \in F[t]$ with $\gcd(a, b) = 1$ and $p \nmid ab$. We call the integer m the *order* of f at p , denoted by $\nu_p(f)$. Conventionally, we set $\nu_p(0) = +\infty$.

Let δ be a derivation on F . Then we can uniquely extend δ to the field $F(t)$ by fixing the value $\delta(t) \in F(t)$. We say that $(F(t), \delta)$ is a *monomial extension* of (F, δ) if t is transcendental over F and $\delta(t) \in F[t]$. We now recall some terminologies for monomial extensions (see Definitions 3.4.2 and 3.5.2 in [8]).

DEFINITION 2.12. Let $(F(t), \delta)$ be a monomial extension of the differential field (F, δ) . A polynomial $p \in F[t]$ is said to be *normal* if $\gcd(p, \delta(p)) = 1$ and it is said to be *special* if $\gcd(p, \delta(p)) = p$. A rational function $f \in F(t)$ is said to be *simple* if the denominator of f is normal and said to be *reduced* if the denominator of f is special.

EXAMPLE 2.13. Let $F = \mathbb{C}$ and $t = x$. If $\delta = d/dx$, then normal polynomials in $F[t]$ are squarefree polynomials and special polynomials are just constants in \mathbb{C} . If $\delta = x \cdot d/dx$, then for any $n \in \mathbb{N}$, x^n is a special polynomial. In fact, we can show that these are the only possible special polynomials.

PROPOSITION 2.14. *Let $(F(t), \delta)$ be a monomial extension of (F, δ) and $f, g \in F(t)$. For any irreducible normal polynomial $p \in F[t]$, we have*

- (i) $v_p(fg) = v_p(f) + v_p(g)$.
- (ii) $v_p(f + g) \geq \min\{v_p(f), v_p(g)\}$ and equality holds if $v_p(f) \neq v_p(g)$.
- (iii) $v_p(\delta(f)) = v_p(f) - 1$ if $v_p(f) \neq 0$, $v_p(\delta(f)) \geq 0$ if $v_p(f) = 0$.
In particular, for any $i \in \mathbb{N}$, $v_p(\delta^i(f)) = v_p(f) - i$ if $v_p(f) < 0$.

PROOF. See Lemma 4.1.1 and Theorem 4.4.2 in [8, Chapter 4]. ■

COROLLARY 2.15. *Let $(F(t), \delta)$ be a monomial extension of (F, δ) and $f \in F(t)$. If f is stable in $(F(t), \delta)$, then $v_p(f) \geq 0$ for any irreducible normal polynomial $p \in F[t]$, i.e., all of the factors of the denominator of f are special polynomials.*

PROOF. Suppose that there exists some irreducible normal polynomial $p \in F[t]$ such that $v_p(f) < 0$. Since f is stable in $(F(t), \delta)$, there exists an element $g_i \in F(t)$ such that $f = \delta^i(g_i)$ for each $i \in \mathbb{N}$. Let $n = -v_p(f)$. By Proposition 2.14 (iii) and the equality $f = \delta^n(g_n)$, we have $v_p(g_n) < 0$ and $v_p(\delta^n(g_n)) = v_p(g_n) - n < -n$, which contradicts with the fact that $v_p(\delta^n(g_n)) = v_p(f) = -n$. ■

3 STABLE ELEMENTARY FUNCTIONS

We now study the stability problem on elementary functions by looking at the classical integration problem of elementary functions through a dynamical lens.

We first recall the differential-algebraic formulation of elementary functions and their integration problems following the presentation in [8].

DEFINITION 3.1. *Let (K, δ) be a differential extension of (k, δ) and $t \in K$. We say that t is elementary over k if one of the following conditions holds:*

- (i) t is algebraic over k , i.e., there exists a polynomial $P \in k[X] \setminus K$ such that $P(t) = 0$;
- (ii) t is exponential over k , i.e., $t \neq 0$ and there exists an element $a \in k$ such that $\delta(t)/t = \delta(a)$, where we write symbolically $t = \exp(a)$;
- (iii) t is logarithmic over k , i.e., there exists an element $a \in k \setminus \{0\}$ such that $\delta(t) = \delta(a)/a$, where we write symbolically $t = \log(a)$.

K is an elementary extension of k if there are $t_1, \dots, t_n \in K$ such that $K = k(t_1, \dots, t_n)$ and t_i is elementary over $k(t_1, \dots, t_{i-1})$ for $i \in \{1, \dots, n\}$. An element $f \in k$ is said to be elementary integrable over k if there exists an elementary extension K of k and $g \in K$ such that $f = \delta(g)$. An elementary function is an element of any elementary extension of the field $(\mathbb{C}(x), d/dx)$.

The classical problem of integration in finite terms is deciding whether a given elementary function is elementary integrable over $\mathbb{C}(x)$ or not. For example, elementary functions

$$\exp(x^2), \quad \frac{\exp(x)}{x}, \quad \frac{\log(x)}{x-1}, \quad \frac{1}{\sqrt{x(x-1)(x-2)}}$$

are not elementary integrable over $\mathbb{C}(x)$. Liouville's theorem is the fundamental principle for elementary integration [8, Chap. 5.5]. For later use, we recall this theorem in the case when the constant field

is algebraically closed since we will study elementary functions over $\mathbb{C}(x)$.

THEOREM 3.2 (LIOUVILLE'S THEOREM). *Let (K, δ) be a differential field with its constant subfield being algebraically closed and $f \in K$. If there exist an elementary extension (E, δ) of (K, δ) with $C_E = C_K$ and $g \in E$ such that $f = \delta(g)$, then there exist $g \in K$, $h_1, \dots, h_n \in K \setminus \{0\}$ and $c_1, \dots, c_n \in C_K$ such that*

$$f = \delta(g) + \sum_{i=1}^n c_i \frac{\delta(h_i)}{h_i}.$$

DEFINITION 3.3. *Let (K, δ) be a differential field and $f \in K$. We say that f is stable over the elementary extensions of K if there exists a sequence $\{g_i\}_{i \in \mathbb{N}}$ such that for all $i \in \mathbb{N}$, g_i is an element of some elementary extension of K and $f = \delta^i(g_i)$. An elementary function that is stable over the elementary extensions of $\mathbb{C}(x)$ is called a stable elementary function.*

LEMMA 3.4. *Let (F, δ) be a regular differential field with $\delta(x) = 1$ and C_F being algebraically closed. Let $(F(t), \delta)$ be a monomial extension of (F, δ) with $C_{F(t)} = C_F$. If $p \in F[t] \setminus F$ is a normal polynomial, then $T = \log(p)$ is not elementary integrable over $F(t)$.*

PROOF. Suppose that T is elementary integrable over $F(t)$. Then Theorem 5.8.2 in [8] implies that there are $c \in C_F$ and $a, b \in F(t)$ such that $T = \delta(cT^2 + aT) + b$ and b is elementary integrable over $F(t)$. By equating the coefficients in T , we get

$$1 = 2c \frac{\delta(p)}{p} + \delta(a) \quad \text{and} \quad a \frac{\delta(p)}{p} + b = 0$$

Since p is normal, Proposition 2.14 (iii) implies that $c = 0$. Then $a = x + \lambda$ for some $\lambda \in C_F$ and $b = -(x + \lambda)\delta(p)/p$. By Lemma 5.6.2 in [8], we get that $x + \lambda$ must be a constant since b is elementary integrable. This is a contradiction. ■

As a special case of the above lemma, we have that $\log(\log(x))$ is not elementary integrable over $\mathbb{C}(x)$.

We now consider the stability problem on elementary functions.

PROBLEM 3.5. *For a given elementary function f over $\mathbb{C}(x)$, decide whether f is stable or not.*

By Lemma 2.10, an elementary function f is stable if and only if for all $i \in \mathbb{N}$, $x^i f$ is elementary integrable over $\mathbb{C}(x)$. So the stability problem can be viewed as a parametrized version of the integration problem. For this moment, we are far from having a complete solution to the above problem. In the rest of this section, we will study the problem on three special families of elementary functions.

3.1 Stable rational functions

It is well-known that all rational functions are elementary integrable over $(\mathbb{C}(x), d/dx)$ since their anti-derivatives are linear combinations of rational functions and logarithmic functions over $\mathbb{C}(x)$.

THEOREM 3.6. *Let f be a rational function in $\mathbb{C}(x)$. Then*

- (i) f is a stable elementary function if $\delta = d/dx$;
- (ii) If $\delta = d/dx$, then f is stable in $(\mathbb{C}(x), \delta)$ if and only if f is a polynomial in $\mathbb{C}[x]$;

(iii) If $\delta = x \cdot d/dx$, then f is stable in $(\mathbb{C}(x), \delta)$ if and only if f is a Laurent polynomial in $\mathbb{C}[x, x^{-1}]$ that is not a nonzero constant.

PROOF. (i) By Lemma 2.10, f is stable in the elementary extensions of $\mathbb{C}(x)$ if and only if for all $i \in \mathbb{N}$, $x^i f$ is elementary integrable. The latter is always true for rational functions.

(ii) Since the differential field $(\mathbb{C}(x), d/dx)$ is regular, all constants in \mathbb{C} are stable by Proposition 2.8. Since 1 is a constant, we have that all polynomials in $\mathbb{C}[x]$ are stable by Theorem 2.4. To show the sufficiency, we assume that $f \in \mathbb{C}(x)$ is stable. By Corollary 2.15, the denominator of f has only special-polynomial factors. Since in the monomial extension $(\mathbb{C}(x), d/dx)$ of $(\mathbb{C}, d/dx)$, the only possible special polynomials are constants in \mathbb{C} . So f must be a polynomial in $\mathbb{C}[x]$.

(iii) By Theorem 5.1.2 in [8], the only possible special polynomials in the monomial extension $(\mathbb{C}(x), x \cdot d/dx)$ of $(\mathbb{C}, x \cdot d/dx)$ are of the form x^i with $i \in \mathbb{N}$. If f is stable in $(\mathbb{C}(x), x \cdot d/dx)$, then f must be a Laurent polynomial in x by Corollary 2.15. Any nonzero constant $c \in \mathbb{C}$ is not integrable (with respect to $x \cdot d/dx$) in $\mathbb{C}(x)$ since $c/x \neq d/dx(g)$ for any $g \in \mathbb{C}(x)$. So f is also not a nonzero constant. The necessity follows from the fact that x^i is stable in $(\mathbb{C}(x), x \cdot d/dx)$ for any nonzero $i \in \mathbb{Z}$ since $x^i = (x \cdot d/dx)^m(x^i/i^m)$ for all $m \in \mathbb{N}$. ■

REMARK 3.7. Another way to show the second assertion in the above theorem is using Theorem 2.11. Suppose that $f \in \mathbb{C}(x)$ is stable and it is not a polynomial. Then $f = P/Q$ for some $P, Q \in \mathbb{C}[x]$ with $Q \notin \mathbb{C}$ and $\gcd(P, Q) = 1$. Since Q is not a constant, Q has at least one root in \mathbb{C} , say α . Let $R = Q/(x - \alpha) \in \mathbb{C}[x]$. By Theorem 2.11, the product $Rf = P/(x - \alpha)$ is also stable in $\mathbb{C}(x)$, which leads to a contradiction since Rf is not integrable in $\mathbb{C}(x)$.

3.2 Stable logarithmic functions

In calculus, we have the following formula [17, p. 238]

$$\int x^n \log(x)^m dx = \frac{x^{n+1}}{m+1} \sum_{k=0}^n (-1)^k \frac{(m+1)!}{(m-k)!} \frac{(\log(x))^{m-k}}{(n+1)^{k+1}} \quad (3.1)$$

for any $m \in \mathbb{N}$ and $n \in \mathbb{Z}$ with $n \neq -1$. This implies that $\log(x)^m$ is a stable elementary functions. One of classical results on the integration of logarithmic functions is as follows.

THEOREM 3.8 (LIOUVILLE-HARDY THEOREM). Let $f \in \mathbb{C}(x)$, then $f \cdot \log(x)$ is elementary integrable over $\mathbb{C}(x)$ if and only if

$$f = \frac{c}{x} + \frac{dg}{dx}$$

for some $c \in \mathbb{C}$ and $g \in \mathbb{C}(x)$.

PROOF. See [19, p. 60] or [28]. ■

Let (K, δ) be a differential field and t be a logarithmic monomial over K , i.e., t is transcendental over K and $\delta(t) = \delta(a)/a$ for some $a \in K \setminus \{0\}$. We further assume that $C_{K(t)} = C_K$. Symbolically, we write $t = \log(a)$. In $K[t]$, all special polynomials are constants by [8, Theorem 5.1.1]. So there is no proper reduced rational function in $K(t)$. For a given $f \in K(t)$, the integration procedure in [8, Chap. 5]

can be summarized as follows. First, applying the Hermite reduction to f yields the decomposition

$$f = \delta(g) + \frac{a}{b},$$

where $g \in K(t)$ and $a, b \in K[t]$ with b being normal. So this step reduces the integrability problem to that of simple rational functions in $K(t)$. By the residue criterion [8, Theorem 5.6.1], there exist $c_1, \dots, c_n \in C_K$ and normal polynomials $b_1, \dots, b_n \in K[t]$ such that $f - \sum_{i=1}^n c_i \delta(b_i)/b_i \in K[t]$ if f is elementary integrable over $K(t)$. If we detect that f is not elementary integrable at this step, we can stop. Otherwise, it remains to consider the integrability problem on polynomials in $K[t]$. For a polynomial $p \in K[t]$, there exists $q \in K[t]$ such that $p - \delta(q) \in K$ if p is elementary integrable over $K(t)$ by [8, Theorem 5.8.1]. So we either detect the non-integrability or reduce the problem from $K(t)$ to K . Then we proceed recursively.

In the rest of this subsection, we specialize to the case in which $K = \mathbb{C}(x)$ and $\delta = d/dx$ and present a stable version of Theorem 3.8.

THEOREM 3.9. Let $T = f \cdot \log(x)$ with $f \in \mathbb{C}(x)$. Then T is stable over the elementary extensions of $\mathbb{C}(x)$ if and only if $f \in \mathbb{C}[x, x^{-1}]$.

PROOF. To show the necessity, we suppose that T is stable over the elementary extension of $\mathbb{C}(x)$. By the Liouville-Hardy theorem, we have $f = c_1/x + \delta(g_1)$ for some $c_1 \in \mathbb{C}$ and $g_1 \in \mathbb{C}(x)$, which implies that

$$f \log(x) = \delta \left(\frac{c_1}{2} (\log(x))^2 + g_1 \log(x) \right) - \frac{g_1}{x}.$$

Since $(\log(x))^2$ and all rational functions are stable, we have that $g_1 \log(x)$ is also stable. Applying the Liouville-Hardy theorem to $g_1 \log(x)$ yields $g_1 = c_2/x + \delta(g_2)$ for some $c_2 \in \mathbb{C}$ and $g_2 \in \mathbb{C}(x)$. Iterating this process, we obtain two sequences $\{c_i\}_{i \in \mathbb{N}}$ in \mathbb{C} and $\{g_i\}_{i \in \mathbb{N}}$ in $\mathbb{C}(x)$ with $g_0 = f$ and $g_i = c_{i+1}/x + \delta(g_{i+1})$ for all $i \in \mathbb{N}$. If the denominator of f has a root other than zero, so do the g_i 's. Choosing sufficiently large i yields a contradiction by looking at the order at this root. So f is a Laurent polynomial in x . For the necessity, we only need to show that $x^m \log(x)$ is stable for any $m \in \mathbb{Z}$. By the formula (3.1), it suffices to show that $\log(x)/x$ is elementary integrable. This is true since $\log(x)/x = \delta(\log(x)^2/2)$. ■

3.3 Stable exponential functions

Throughout this part, let δ be the usual derivation d/dx on $\mathbb{C}(x)$ and its extensions. The non-elementary integrability of $\exp(x^2)$ is derived from the following theorem by Liouville in [24] (see [41, p. 971] for its proof).

THEOREM 3.10. Let $f, g \in \mathbb{C}(x)$ with $g \notin \mathbb{C}$ and $t = f \cdot \exp(g)$. Then t is elementary integrable over $\mathbb{C}(x)$ if and only if there exists $h \in \mathbb{C}(x)$ such that $f = \delta(h) + h \cdot \delta(g)$.

A rational function $f = a/b$ with $a, b \in \mathbb{C}[x]$ and $\gcd(a, b) = 1$ is said to be *differential-reduced* (with respect to δ) if

$$\gcd(b, a - i\delta(b)) = 1 \quad \text{for all } i \in \mathbb{Z}.$$

We recall some basic properties of differential-reduced rational functions from [6, 13].

PROPOSITION 3.11. Let $f = a/b \in \mathbb{C}(x)$ be such that $a, b \in \mathbb{C}[x]$ and $\gcd(a, b) = 1$. Then

(i) $\delta(f)$ is always differential-reduced;

- (ii) $f + m\delta(b)/b$ is differential-reduced for any $m \in \mathbb{Z}$ if f is differential-reduced;
- (iii) if f is differential-reduced and $b\delta(g) + ag \in \mathbb{C}[x]$, then $g \in \mathbb{C}[x]$.

PROOF. The first two assertions follows from Lemma 2 in [13] saying that f is differential-reduced if and only if none of its residues at simple poles is an integer. The third one is Lemma 6 in [6]. ■

LEMMA 3.12. *Let $g \in \mathbb{C}(x)$ and $f = \delta(g) = a/b$ with $a, b \in \mathbb{C}[x]$ and $\gcd(a, b) = 1$. Then $t = P \cdot b^m \cdot \exp(g)$ with $m \in \mathbb{N}$ and $P \in \mathbb{C}[x] \setminus \{0\}$ is elementary integrable over $\mathbb{C}(x)$ if and only if there exists $Q \in \mathbb{C}[x]$ such that $t = \delta(Q \cdot b^{m+1} \cdot \exp(g))$ and*

$$\deg_x(Q) = \begin{cases} \deg_x(P) - \deg_x(a), & \text{if } \deg_x(a) \geq \deg_x(b); \\ \deg_x(P) - \deg_x(b) + 1, & \text{if } \deg_x(a) < \deg_x(b) - 1. \end{cases}$$

PROOF. Suppose that t is elementary integrable over $\mathbb{C}(x)$. Then Theorem 3.10 implies that there exists $h \in \mathbb{C}(x)$ such that

$$Pb^m = \delta(h) + f \cdot h.$$

Write $h = Q \cdot b^{m+1}$ with $Q \in \mathbb{C}(x)$. Then

$$P = b\delta(Q) + (a + (m+1)\delta(b))Q.$$

We first show that $Q \in \mathbb{C}[x]$. Since $f = \delta(g) = a/b$, Proposition 3.11 (i) and (ii) imply that both f and $(a + (m+1)\delta(b))/b$ are differential-reduced. Then we get $Q \in \mathbb{C}[x]$ by Proposition 3.11 (iii). We next estimate $\deg_x(Q)$. Since $f = \delta(g)$, we have either $\deg_x(a) \geq \deg_x(b)$ or $\deg_x(a) < \deg_x(b) - 1$ by Theorem 4.4.4 in [8], i.e., $\deg_x(a) \neq \deg_x(b) - 1$. If $\deg_x(a) \geq \deg_x(b)$, then $\deg_x(P) = \deg_x(a) + \deg_x(Q)$. Hence $\deg_x(Q) = \deg_x(P) - \deg_x(a)$. If $\deg_x(a)$ is less than $\deg_x(b) - 1$, then the leading monomial of

$$b\delta(Q) + (a + (m+1)\delta(b))Q$$

is equal to

$$\text{lc}(Q) \cdot \text{lc}(b) \cdot (\deg_x(Q) + (m+1)\deg_x(b))x^{\deg_x(b) + \deg_x(Q) - 1},$$

which implies that $\deg_x(Q) = \deg_x(P) - \deg_x(b) + 1$. ■

We now present a stable version of Theorem 3.10.

THEOREM 3.13. *Let $f, g \in \mathbb{C}(x)$ be such that $g \notin \mathbb{C}$ and $t = f \cdot \exp(g)$. Then t is a stable elementary function if and only if $f \in \mathbb{C}[x]$ and $g = \lambda x + \mu$ for some $\lambda, \mu \in \mathbb{C}$ with $\lambda \neq 0$.*

PROOF. To see the necessity, note that the function $\exp(\lambda x + \mu)$ is stable since for any $i \in \mathbb{N}$,

$$\exp(\lambda x + \mu) = \delta^i \left(\frac{1}{\lambda^i} \exp(\lambda x + \mu) \right).$$

By Theorem 2.11, $f \cdot \exp(\lambda x + \mu)$ is stable for any $f \in \mathbb{C}[x]$. For the sufficiency, we assume that t is a stable elementary function. Since $g \notin \mathbb{C}$, we have $\delta(g) \neq 0$. We first claim that $\delta(g)$ must be a constant. Suppose that this is not true. Let d be the denominator of f . Then $t_1 = d \cdot t = p \cdot \exp(g)$ for some $p \in \mathbb{C}[x]$ and it is also stable by Theorem 2.11. Since t_1 is elementary integrable over $\mathbb{C}(x)$, Theorem 3.10 implies that there exists $h \in \mathbb{C}(x)$ such that

$$p = \delta(h) + \delta(g) \cdot h.$$

Write $\delta(g) = a/b$ with $a, b \in \mathbb{C}[x]$ and $\gcd(a, b) = 1$. By Lemma 3.12, there exists $Q_1 \in \mathbb{C}[x]$ such that $h = bQ_1$ and

$$\deg_x(Q_1) = \begin{cases} \deg_x(p) - \deg_x(a), & \text{if } \deg_x(a) \geq \deg_x(b); \\ \deg_x(p) - \deg_x(b) + 1, & \text{if } \deg_x(a) < \deg_x(b) - 1. \end{cases}$$

Notice that $\deg_x(Q_1) < \deg_x(p)$ since $\delta(g)$ is not a constant. Since t_1 is stable and $t_1 = \delta(Q_1 b \exp(g))$, we have $t_2 = Q_1 b \exp(g)$ is also stable. Following the same argument as above, we have $t_2 = \delta(Q_2 \cdot b^2 \exp(g))$ for some $Q_2 \in \mathbb{C}[x]$ with $\deg_x(Q_2) < \deg_x(Q_1)$. Repeating this process, we obtain a sequence of polynomials $\{Q_i\}_{i \in \mathbb{N}}$ with $Q_0 = p$ and $\deg_x(Q_{i+1}) < \deg_x(Q_i)$. This is a contradiction. Thus, $\delta(g)$ is a nonzero constant and then $g = \lambda x + \mu$ for some $\lambda, \mu \in \mathbb{C}$ with $\lambda \neq 0$. It remains to show that $f \in \mathbb{C}[x]$. Suppose that $f \in \mathbb{C}(x)$ not a polynomial. Then $f = P/Q$ for some $P, Q \in \mathbb{C}[x]$ with $Q \notin \mathbb{C}$ and $\gcd(P, Q) = 1$. Since Q is not a constant, Q has at least one root in \mathbb{C} , say α . Let $R = Q/(x - \alpha) \in \mathbb{C}[x]$. Since $f \exp(g)$ is stable, so is $R \cdot f \exp(g) = P \exp(g)/(x - \alpha)$ by Theorem 2.11. Now again Theorem 3.10 implies that there exists $h \in \mathbb{C}(x)$ such that

$$\frac{P}{x - \alpha} = \delta(h) + \lambda \cdot h.$$

By estimating the valuation at $x - \alpha$, we have $v_{x-\alpha}(P/(x - \alpha)) = -1$ but $v_{x-\alpha}(\delta(h) + \lambda \cdot h)$ is either non-negative or strictly less than -1 , which leads to a contradiction. ■

4 STABLE D-FINITE POWER SERIES

The notion of D-finite power series was first introduced by Stanley [47] in 1980 and studied extensively in [25, 48, 52]. These series like algebraic numbers can be algorithmically manipulated via its defining linear differential equations [1, 44]. We will study the stability problem on D-finite power series.

Let k be a field of characteristic zero and let $k[[x]]$ be the ring of formal power series in x over k and let δ denote the derivation with $\delta(x) = 1$ and $\delta(a) = 0$ for all $a \in k$. The quotient field of $k[[x]]$ is called the field of formal Laurent series, denoted by $k((x))$, which includes $k(x)$ as its subfield. Let R be a ring and $\sigma : R \rightarrow R$ be an isomorphism on R . We call the pair (R, σ) a *difference ring*. Over a difference ring, we have the polynomial ring $R\langle S \rangle$ in which the addition is defined coefficient-wise in the indeterminate S and the multiplication satisfies the rule:

$$S \cdot r = \sigma(r) \cdot S \quad \text{for all } r \in R.$$

In particular, we call $k[x]\langle S \rangle$ the ring of linear recurrence operators with polynomial coefficients. For a sequence $a_n : \mathbb{N} \rightarrow k$, we define an action of an operator $P = \sum_{i=0}^d p_i S^i \in k[x]\langle S \rangle$ on a_n by $P(a_n) = \sum_{i=0}^d p_i(n) a_{n+i}$. For an operator $L = \sum_{i=0}^d \ell_i D^i \in k[x]\langle D \rangle$ with $\ell_d \neq 0$, we call d the order of L , denoted by $\text{ord}(L)$ and $\max\{\deg_x(\ell_0), \dots, \deg_x(\ell_d)\}$ the degree of L , denoted by $\deg(L)$. We define the order and the degree of an operator $P \in k[x]\langle S \rangle$ in a similar way.

DEFINITION 4.1. *A power series $f = \sum_{n \geq 0} a_n x^n \in k[[x]]$ is said to be D-finite over $k(x)$ if there exists a non-zero operator $L \in k[x]\langle D \rangle$ such that $L(f) = 0$. Such an operator L is called an annihilator for f . A sequence $a_n : \mathbb{N} \rightarrow K$ is said to be P-recursive if there exists a non-zero operator $P \in k[x]\langle S \rangle$ such that $P(a_n) = 0$. We also call such an operator P an annihilator for a_n .*

The following theorem summarizes some fundamental properties of D-finite power series and P-recursive sequences.

THEOREM 4.2. *Let $f = \sum_{n \geq 0} a_n x^n \in k[[x]]$. Then*

- (i) *f is D-finite if and only if its coefficient sequence a_n is P-recursive;*
- (ii) *if a_n has an annihilator in $k[x]\langle S \rangle$ of order r and degree d , then f has an annihilator in $k[x]\langle D \rangle$ of order at most d and degree at most $r + d$;*
- (iii) *if both a_n and b_n are P-recursive with annihilators A and B in $k[x]\langle S \rangle$, then $a_n b_n$ is also P-recursive with an annihilator of order at most $\text{ord}(A)\text{ord}(B)$ and degree at most*

$$2 \max\{\text{deg}(A), \text{deg}(B)\} \text{ord}(A)^2 \text{ord}(B)^2.$$

PROOF. For the proofs, see [22, p. 149] and [21, Theorem 8]. ■

As a high-order generalization of Gosper's algorithm [16] and its differential analogue [4], Abramov and van Hoeij studied the integration problem on solutions of linear functional equations [2, 3]. In the following, we let $k = \mathbb{C}$ and $\delta = d/dx$ so that we can talk about generalized series solutions of linear differential equations.

PROBLEM 4.3. *Given an operator $L \in k(x)\langle D \rangle$, find a minimal-order operator $\tilde{L} \in k(x)\langle D \rangle$ such that the derivatives of the solutions of \tilde{L} are the solutions of L . We call \tilde{L} an integral of L , denoted by $\text{int}(L)$.*

Note that \tilde{L} is unique up to a factor in K and the order of \tilde{L} is greater than that of L by at most one. If $\text{ord}(\tilde{L}) = \text{ord}(L)$, Abramov and van Hoeij proved that there exist $P \in k(x)\langle D \rangle$ with $\text{ord}(P) < \text{ord}(L)$ and $r \in k(x)$ such that $D \cdot P + r \cdot L = 1$. In this case, a solution f of L has an anti-derivative of the form $P(f)$. For a power series $f = \sum_{n \geq 0} a_n x^n \in k[[x]]$, we call the series $\sum_{n \geq 1} \frac{a_{n-1}}{n} x^n$ a formal integral of f , denoted by $\text{int}(f)$.

DEFINITION 4.4. *A D-finite power series f is said to be stable if there exists a sequence $\{g_i\}_{i \in \mathbb{N}}$ in $k[[x]]$ such that $g_0 = f$, $g_i = \delta(g_{i+1})$ and all of the g_i 's have annihilators of the same order. It is said to be eventually stable if there exists $m \in \mathbb{N}$ such that $\text{int}^m(f)$ is stable. An operator $L \in k(x)\langle D \rangle$ is said to be stable if $\text{ord}(\text{int}^i(L)) = \text{ord}(L)$ for all $i \in \mathbb{N}$ and be eventually stable if there exists $m \in \mathbb{N}$ such that $\text{int}^m(L)$ is stable.*

The following result was first discovered by Guo and then proved by the author in 2020. For more interesting stable power series, see Guo's thesis [18].

THEOREM 4.5. *Any D-finite power series is eventually stable.*

PROOF. Let $f = \sum_{n \geq 0} a_n x^n \in k[[x]]$ be D-finite and $L \in k[x]\langle D \rangle$ be the minimal annihilator for f . By Theorem 4.2 (i), a_n is P-recursive and so has a minimal annihilator $P \in k(x)\langle S \rangle$. Let

$$g_i = \sum_{n \geq i} \frac{a_{n-i}}{n(n-1) \cdots (n-i+1)} x^n.$$

Then $f = \delta^i(g_i)$. We claim that the order of minimal annihilators for g_i is bounded. By Theorem 4.2 (ii), it suffices to show that the degrees of the minimal annihilators of the coefficient sequences $b_{n,i} = a_{n-i}/(n(n-1) \cdots (n-i+1))$ are bounded. Note that a_{n-i} has the same annihilator as a_n and the minimal annihilators of the sequences $t_i = 1/(n(n-1) \cdots (n-i+1))$ are $(n+1)S - (n-i+1)$,

whose order and degree are independent of i . Then Theorem 4.2 (iii) implies that the degrees of the minimal annihilators for $b_{n,i}$ are at most $2 \max\{1, \text{deg}(P)\} \text{ord}(P)^2$. Then there exists $m \in \mathbb{N}$ such that the formal integral $\text{int}^m(f)$ is stable. ■

5 CONCLUSION AND FUTURE WORK

This paper presents some initial results towards a deep connection between dynamics and differential algebra with the focus on stability problems in symbolic integration. This is just a first try and more general cases are waiting for further studying in this direction.

To conclude this paper, we propose some problems for future work. The first problem is characterizing all possible algebraic functions that are stable in the differential field $(\overline{\mathbb{C}(x)}, d/dx)$. The typical stable family of algebraic functions is $(x-c)^r$ with $c \in \mathbb{C}$ and $r \in \mathbb{Q} \setminus \{-1, -2, \dots\}$. We conjecture that an algebraic function is stable in $(\overline{\mathbb{C}(x)}, d/dx)$ if and only if it is of the form

$$\sum_{i=1}^n p_i \cdot (x - c_i)^{r_i},$$

where $p_i \in \mathbb{C}[x]$, $c_i \in \mathbb{C}$ and $r_i \in \mathbb{Q} \setminus \{-1, -2, \dots\}$. The second problem is formulating a stable version of Liouville's theorem that describes the structure of elementary functions that are elementary integrable. This will be crucial for developing a Risch-type algorithm for detecting whether an elementary function is stable or not. The third problem is studying stability problems in symbolic summation. For this moment, we have some parallel results in this direction which will be included in a forthcoming paper. A special case of this problem is to characterize all possible stable hypergeometric terms with respect to the difference operator by thinking Gosper's algorithm dynamically. The last problem is related to a classical open problem in differential algebra asked by Rubel in [43]. For a D-finite power series $f = \sum_{n \geq 0} a_n x^n \in k[[x]]$, Rubel conjectured that the set $\{n \in \mathbb{N} \mid a_n = 0\}$ is a Skolem set, i.e., a union of finitely many arithmetic progressions. A given elementary function $f(x)$ over $\mathbb{C}(x)$ is generically not stable, but we conjectured that the set

$$\{i \in \mathbb{N} \mid x^i f(x) \text{ is elementary integrable over } \mathbb{C}(x)\}$$

is also a Skolem set. For instance, $\exp(x^2)$ is not elementary integrable, but for odd $i \in \mathbb{N}$, $x^i \exp(x^2)$ is elementary integrable.

Acknowledgement. I would like to thank Ruyong Feng and Umberto Zannier for many discussions during the formation of the conjecture for stable algebraic functions and Ze-wang Guo for sharing his discoveries and many interesting examples on the exact integration of D-finite functions. I am also very grateful to the two anonymous referees for their constructive comments and to Hao Du, Lixin Du, Hanqian Fang, Pingchuan Ma, and Chaochao Zhu for reading the draft and sending me their lists of typos.

REFERENCES

- [1] Sergei A. Abramov, Ha Quang Le, and Ziming Li. Univariate Ore polynomial rings in computer algebra. *J. of Mathematical Sci.*, 131(5):5885–5903, 2005.
- [2] Sergei A. Abramov and Mark van Hoeij. A method for the integration of solutions of Ore equations. In *Proceedings of the 1997 International Symposium on Symbolic and Algebraic Computation (ISSAC'97)*, pages 172–175. ACM, New York, 1997.

- [3] Sergei A. Abramov and Mark van Hoeij. Integration of solutions of linear functional equations. *Integral Transform. Spec. Funct.*, 8(1-2):3–12, 1999.
- [4] Gert Almkvist and Doron Zeilberger. The method of differentiating under the integral sign. *J. Symbolic Comput.*, 10:571–591, 1990.
- [5] Alan F. Beardon. *Iteration of rational functions*, volume 132 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1991. Complex analytic dynamical systems.
- [6] Alin Bostan, Shaoshi Chen, Frédéric Chyzak, Ziming Li, and Guoce Xin. Hermite reduction and creative telescoping for hyperexponential functions. In *Proceedings of the 2013 International Symposium on Symbolic and Algebraic Computation (ISSAC '13)*, pages 77–84, New York, NY, USA, 2013. ACM.
- [7] Manuel Bronstein. Integration of elementary functions. *J. Symbolic Comput.*, 9(2):117–173, 1990.
- [8] Manuel Bronstein. *Symbolic Integration I: Transcendental Functions*, volume 1 of *Algorithms and Computation in Mathematics*. Springer-Verlag, Berlin, second edition, 2005.
- [9] Manuel Bronstein and Marko Petkovšek. An introduction to pseudo-linear algebra. *Theoret. Comput. Sci.*, 157:3–33, 1996.
- [10] James H. Davenport. Complexity of integration, special values, and recent developments. In *Mathematical software—ICMS 2016*, volume 9725 of *Lecture Notes in Comput. Sci.*, pages 485–491. Springer, [Cham], 2016.
- [11] James Harold Davenport. *On the Integration of Algebraic Functions*, volume 102 of *Lecture Notes in Computer Science*. Springer-Verlag, Berlin, 1981.
- [12] Keith O. Geddes, Stephanie R. Czapor, and George Labahn. *Algorithms for Computer Algebra*. Kluwer Academic Publishers, Boston, MA, 1992.
- [13] Keith O. Geddes, Ha Quang Le, and Ziming Li. Differential rational normal forms and a reduction algorithm for hyperexponential functions. In *Proceedings of the 2004 International Symposium on Symbolic and Algebraic Computation (ISSAC'04)*, pages 183–190, New York, USA, 2004. ACM.
- [14] Jürgen Gerhard. Fast modular algorithms for squarefree factorization and Hermite integration. *Appl. Algebra Engrg. Comm. Comput.*, 11(3):203–226, 2001.
- [15] Eddy Godelle. The stable set of a self-map. *Adv. in Appl. Math.*, 45(3):438–448, 2010.
- [16] Ralph William Gosper, Jr. Decision procedure for indefinite hypergeometric summation. *Proc. Nat. Acad. Sci. U.S.A.*, 75(1):40–42, 1978.
- [17] I. S. Gradshteyn and I. M. Ryzhik. *Table of integrals, series, and products*. Elsevier/Academic Press, Amsterdam, seventh edition, 2007. Translated from the Russian.
- [18] Zewang Guo. Integration of Single Variable D-finite Functions, Bachelor's Thesis (Beihang University), 2020 (in Chinese).
- [19] G. H. Hardy. *The integration of functions of a single variable*. Cambridge Tracts in Mathematics and Mathematical Physics, No. 2. Hafner Publishing Co., New York, 1971. Reprint of the second edition, 1916.
- [20] Toni Kasper. Integration in finite terms: the Liouville theory. *Math. Mag.*, 53(4):195–201, 1980.
- [21] Manuel Kauers. Bounds for D-finite closure properties. In *Proceedings of the 2014 International Symposium on Symbolic and Algebraic Computation (ISSAC'14)*, pages 288–295. ACM, New York, 2014.
- [22] Manuel Kauers and Peter Paule. *The Concrete Tetrahedron: Symbolic sums, Recurrence equations, Generating functions, Asymptotic estimates..* Texts and Monographs in Symbolic Computation. SpringerWienNewYork, Vienna, 2011.
- [23] Askold Khovanskii. *Topological Galois theory*. Springer Monographs in Mathematics. Springer, Heidelberg, 2014.
- [24] Joseph Liouville. Mémoire sur l'intégration d'une classe de fonctions transcendentes. *J. Reine Angew. Math.*, 13:93–118, 1835.
- [25] Leonard M. Lipshitz. D-finite power series. *J. Algebra*, 122(2):353–373, 1989.
- [26] Jesper Lützen. *Joseph Liouville 1809–1882: master of pure and applied mathematics*, volume 15 of *Studies in the History of Mathematics and Physical Sciences*. Springer-Verlag, New York, 1990.
- [27] Elena Anne Marchisotto and Gholam-Ali Zakeri. An invitation to integration in finite terms. *The College Mathematics Journal*, 25(4):295–308, 1994.
- [28] Elena Anne Marchisotto and Gholam-Ali Zakeri. An invitation to integration in finite terms. *The College Mathematics Journal*, 25(4):295–308, 1994.
- [29] David Masser. Integration in elementary terms. *Newsletter London Mathematical Society*, 47:30–36, 2017.
- [30] David Masser and Umberto Zannier. Torsion points, Pell's equation, and integration in elementary terms. *Acta Math.*, 225(2):227–313, 2020.
- [31] Joel Moses. *Symbolic Integration*. PhD Thesis, Massachusetts Institute of Technology, 1968.
- [32] Saradha Natarajan and Ravindranathan Thangadurai. *Pillars of transcendental number theory*. Springer, Singapore, 2020.
- [33] Ivan Niven. *Irrational numbers*. The Carus Mathematical Monographs, No. 11. The Mathematical Association of America. Distributed by John Wiley and Sons, Inc., New York, N.Y., 1956.
- [34] Oystein Ore. Theory of non-commutative polynomials. *Ann. of Math. (2)*, 34(3):480–508, 1933.
- [35] Alexandre Ostrowski. Sur les relations algébriques entre les intégrales indéfinies. *Acta Math.*, 78:315–318, 1946.
- [36] Clemens G. Raab. *Definite Integration in Differential Fields*. PhD thesis, Johannes Kepler Universität Linz, Austria, August 2012.
- [37] Clemens G. Raab. Generalization of Risch's algorithm to special functions. In *Computer algebra in quantum field theory*, Texts Monogr. Symbol. Comput., pages 285–304. Springer, Vienna, 2013.
- [38] Robert H. Risch. The problem of integration in finite terms. *Trans. Amer. Math. Soc.*, 139:167–189, 1969.
- [39] Robert H. Risch. The solution of the problem of integration in finite terms. *Bull. Amer. Math. Soc.*, 76:605–608, 1970.
- [40] Joseph Fels Ritt. *Integration in Finite Terms. Liouville's Theory of Elementary Methods*. Columbia University Press, New York, N. Y., 1948.
- [41] Maxwell Rosenlicht. Integration in finite terms. *Amer. Math. Monthly*, 79:963–972, 1972.
- [42] Michael Rothstein. *Aspects of Symbolic Integration and Simplification of Exponential and Primitive Functions*. PhD Thesis, University of Wisconsin, 1976.
- [43] Lee A. Rubel. Some research problems about algebraic differential equations. *Trans. Amer. Math. Soc.*, 280(1):43–52, 1983.
- [44] Bruno Salvy. Linear differential equations as a data structure. *Found. Comput. Math.*, 19(5):1071–1112, 2019.
- [45] Joseph H. Silverman. *The Arithmetic of Dynamical Systems*, volume 241 of *Graduate Texts in Mathematics*. Springer, New York, 2007.
- [46] James Robert Slagle. *A Heuristic Program that Solves Symbolic Integration Problems in Freshman Calculus, Symbolic Automatic Integrator (SAINT)*. PhD Thesis, Massachusetts Institute of Technology, 1961.
- [47] Richard P. Stanley. Differentiably finite power series. *European J. Combin.*, 1(2):175–188, 1980.
- [48] Richard P. Stanley. *Enumerative combinatorics. Vol. 2*, volume 62 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1999.
- [49] Barry M. Trager. *On the Integration of Algebraic Functions*. PhD Thesis, Massachusetts Institute of Technology, 1984.
- [50] Joachim von zur Gathen and Gerhard Jürgen. *Modern Computer Algebra*. Cambridge University Press, Cambridge, second edition, 2003.
- [51] Umberto Zannier. Elementary integration of differentials in families and conjectures of Pink. In *Proceedings of the International Congress of Mathematicians—Seoul 2014. Vol. II*, pages 531–555. Kyung Moon Sa, Seoul, 2014.
- [52] Doron Zeilberger. A holonomic systems approach to special functions identities. *J. Comput. Appl. Math.*, 32:321–368, 1990.