

ALGORITHMS YIELD UPPER BOUNDS IN DIFFERENTIAL ALGEBRA

WEI LI, ALEXEY OVCHINNIKOV, GLEB POGUDIN, AND THOMAS SCANLON

ABSTRACT. Consider an algorithm computing in a differential field with several commuting derivations such that the only operations it performs with the elements of the field are arithmetic operations, differentiation, and zero testing. We show that, if the algorithm is guaranteed to terminate on every input, then there is a computable upper bound for the size of the output of the algorithm in terms of the input. We also generalize this to algorithms working with models of good enough theories (including for example, difference fields).

We then apply this to differential algebraic geometry to show that there exists a computable uniform upper bound for the number of components of any variety defined by a system of polynomial PDEs. We then use this bound to show the existence of a computable uniform upper bound for the elimination problem in systems of polynomial PDEs with delays.

1. INTRODUCTION

Finding uniform bounds for problems and quantities (e.g., consistency testing or counting of solutions) is one of the central questions in differential algebra. In [26], it was demonstrated that, in commutative algebra, one can show the existence of such bounds as a consequence of theorems about nonstandard extensions of standard algebraic objects. This approach was successfully applied in the differential algebra context in [11] and [8, Section 6] for establishing, for example, the existence of a uniform bound in the differential Nullstellensatz. Furthermore, in [25], the authors used methods of proof theory to extract explicit bounds based on nonstandard existence proofs.

The present paper can be viewed as an alternative approach, in which we derive the existence of a computable uniform bound for an object from the existence of an algorithm for computing the object. More precisely, let T be a complete recursive theory. The most relevant examples for us would be the theory of differentially closed fields in zero characteristic with m commuting derivations and the theory of existentially closed difference fields, others include algebraically closed and real closed fields. Consider an algorithm A performing computations in a model of T that is restricted to using only definable functions when working with elements of the model (for formal definition, we refer to Section 4.1) and required to terminate for every input.

We show that there is a computable upper bound for the size of the output of A in terms of the input size of A . We apply this to the Rosenfeld-Gröbner algorithm [2] that decomposes a solution set of a system of polynomial PDEs into components and is such an algorithm. This allows us to show that there is a uniform upper

2010 *Mathematics Subject Classification*. Primary 12H05, 12H10, 03C10; Secondary 03C60, 03D15.

bound for the number of components of any differential-algebraic variety defined by a system of polynomial PDEs. We also show how this bound for the number of components leads to a uniform upper bound for the elimination problem in systems of polynomial PDEs with delays.

A bound for the number of components of varieties defined by polynomial ODEs appeared in [18], as did a bound for the elimination problem for polynomial ODEs with delays. These bounds are based on the application of the Rosenfeld-Gröbner algorithm, which, if applied in this situation to ODEs, outputs equations whose order does not exceed the order of the input. This allowed to restrict to a finitely generated subring of the ring of differential polynomials and use tools from algebraic geometry. It is non-trivial to generalize this to polynomial PDEs because the orders in the output of the Rosenfeld-Gröbner can be larger than the orders of the input. Another key ingredient in the ODE case to obtain the bound in [18] was an analysis of differential dimension polynomials. A significant difference of our present PDE context with the ordinary case is that these polynomials behave less predictably under projections of varieties (compare [18, Lemma 6.16] and Lemma 6.3). To overcome this difficulty, we use again our bound for the Rosenfeld-Gröbner algorithm.

We believe that our method can also be applied to obtain bounds for other algorithms in differential algebra such as [1, Algorithm 3.6] and for algorithms from other theories, e.g. [7, Algorithm 3] for systems of difference equations. Since the reducibility of a polynomial can be expressed as a first-order existential formula, it seems plausible that the same methods could be applied to other algorithms dealing with difference [5] and differential-difference [6] equations that use factorization because the corresponding theories satisfy the requirements of our approach [14, 17, 23]. However, we leave these for future research.

The paper is organized as follows. Section 2 contains definitions and notation used in Section 3 to state the main results. Bounds for an algorithm working with a model of a theory T are established in Section 4. These results are applied to differential algebra in Section 5. Further applications to delay PDEs are given in Section 6.

2. BASIC NOTIONS AND NOTATION

Definition 2.1 (Differential-difference rings).

- A Δ - σ -ring $(\mathcal{R}, \Delta, \sigma)$ is a commutative ring \mathcal{R} endowed with a finite set $\Delta = \{\partial_1, \dots, \partial_m\}$ of commuting derivations of \mathcal{R} and an endomorphism σ of \mathcal{R} such that, for all i , $\partial_i \sigma = \sigma \partial_i$.
- When \mathcal{R} is additionally a field, it is called a Δ - σ -field.
- If σ is an automorphism of \mathcal{R} , \mathcal{R} is called a Δ - σ^* -ring.
- If $\sigma = \text{id}$, \mathcal{R} is called a Δ -ring or differential ring.
- For a commutative ring R , $\langle F \rangle$ denotes the ideal generated by $F \subset R$ in R .
- For $\Delta = \{\partial_1, \dots, \partial_m\}$, let $\Theta_\Delta = \{\partial_1^{i_1} \cdots \partial_m^{i_m} \mid i_j \geq 0, 1 \leq j \leq m\}$.
- For $\theta = \partial_1^{i_1} \cdots \partial_m^{i_m} \in \Theta_\Delta$, we let $\text{ord } \theta = i_1 + \dots + i_m$. For a non-negative integer B , we denote $\Theta_\Delta(B) := \{\theta \in \Theta_\Delta \mid \text{ord } \theta \leq B\}$.
- For a Δ -ring R , the differential ideal generated by $F \subset R$ in R is denoted by $\langle F \rangle^{(\infty)}$; for a non-negative integer B , we introduce the following ideal of R :

$$\langle F \rangle^{(B)} := \langle \theta(F) \mid \theta \in \Theta_\Delta(B) \rangle.$$

Definition 2.2 (Differential polynomials). Let \mathcal{R} be a Δ -ring. The differential polynomial ring over \mathcal{R} in $\mathbf{y} = y_1, \dots, y_n$ is defined as

$$\mathcal{R}\{\mathbf{y}\}_\Delta := \mathcal{R}[\theta y_k \mid \theta \in \Theta_\Delta; 1 \leq k \leq n].$$

The structure of a Δ -ring is defined by $\partial_i(\theta y_k) := (\partial_i \theta) y_k$ for every $\theta \in \Theta_\Delta$.

Definition 2.3 (Differential-difference polynomials). Let \mathcal{R} be a Δ - σ -ring. The differential-difference polynomial ring over \mathcal{R} in $\mathbf{y} = y_1, \dots, y_n$ is defined as

$$\mathcal{R}[\mathbf{y}_\infty] := \mathcal{R}[\theta \sigma^i y_k \mid \theta \in \Theta_\Delta; i \geq 0; 1 \leq k \leq n].$$

The structure of Δ - σ ring is defined by $\sigma(\theta \sigma^j y_k) := \theta \sigma^{j+1} y_k$ and $\partial_i(\theta \sigma^j y_k) := (\partial_i \theta) \sigma^j y_k$ for every $\theta \in \Theta_\Delta$ and $j \geq 0$.

A Δ - σ -polynomial is an element of $\mathcal{R}[\mathbf{y}_\infty]$. Given $B \in \mathbb{N}$, let $\mathcal{R}[\mathbf{y}_B]$ denote the polynomial ring

$$\mathcal{R}[\theta \sigma^j y_k \mid \theta \in \Theta_\Delta(B); 0 \leq j \leq B; 1 \leq k \leq n].$$

For the notions from logic that we use, see [19, Sections 2.1-2.2].

3. MAIN RESULTS

For clarity, we gather our main results in one section.

Theorem 3.1 (Upper bound for irreducible components for PDEs). *There exists a computable function $\text{Comp}(m, n)$ such that, for every differential field k with a set of m commuting derivations Δ and finite $F \subset k\{y_1, \dots, y_n\}_\Delta$ with $\max\{\text{ord } F, \text{deg } F\} \leq s$, the number of components in the variety defined by $F = 0$ does not exceed $\text{Comp}(m, \max\{n, s\})$.*

Additional details and proof are given in Theorem 5.11.

Theorem 3.2 (Upper bound for elimination in delay PDEs). *For all non-negative integers r, m and s , there exists a computable $B = B(r, m, s)$ such that, for all:*

- *non-negative integers q and t ,*
- *a Δ - σ -field k with $\text{char } k = 0$ and $|\Delta| = m$,*
- *sets of Δ - σ -polynomials $F \subset k[\mathbf{x}_t, \mathbf{y}_s]$, where $\mathbf{x} = x_1, \dots, x_q$, $\mathbf{y} = y_1, \dots, y_r$, and $\text{deg}_{\mathbf{y}} F \leq s$,*

we have

$$\begin{aligned} \langle \sigma^i(F) \mid i \in \mathbb{Z}_{\geq 0} \rangle^{(\infty)} \cap k[\mathbf{x}_\infty] \neq \{0\} \\ \iff \langle \sigma^i(F) \mid i \in [0, B] \rangle^{(B)} \cap k[\mathbf{x}_{B+t}] \neq \{0\}. \end{aligned}$$

The two preceding theorems are proved using our main technical result about algorithms performing computations in complete recursive theories. Stating it precisely requires defining admissible algorithms carefully, so we postpone it until Section 4 and give here a simplified and informal version of the statement.

Theorem 3.3 (Algorithm yields a bound, stated precisely as Theorem 4.5). *There exists a computable function with input*

- *complete recursive theory T ;*
- *an algorithm \mathcal{A} performing computations in a model of T restricted to using only definable functions when working with elements of the model;*
- *positive integer ℓ*

that computes a number N such that for every model M of T and every $\mathbf{a} \in M^\ell$ the size of the output of \mathcal{A} with input \mathbf{a} does not exceed N .

For the application of this to the Rosenfeld-Gröbner algorithm, see Theorem 5.8.

4. BOUNDS FOR THE OUTPUT SIZE OF ALGORITHMS OVER COMPLETE THEORIES

In this section, we will use the formalism of oracle Turing machines [24, § 14.3]. Roughly speaking, an oracle Turing machine is a Turing machine with an extra tape for performing queries to an external oracle. An oracle is not considered to be a part of the machine.

4.1. Setup. To consider an algorithm dealing with elements of a (not necessarily computable) model of a theory T , we will “encapsulate” the elements of the model given to the algorithm into an oracle that allows to perform only first-order operations with them as defined below. For other approaches that could be used to formalizing computations in arbitrary structures, see [9, §1] and [4, §2.2].

Definition 4.1 (T -oracle). Let \mathcal{L} be a language and T be a theory in \mathcal{L} . For elements a_1, \dots, a_ℓ of a model M of T , any oracle that supports the following queries: given a formula $\varphi(x_1, \dots, x_\ell)$, the oracle returns the value $\varphi(a_1, \dots, a_\ell)$ in M , will be denoted by $\mathcal{O}_M(a_1, \dots, a_\ell)$.

Definition 4.2 (Total algorithm over T). An oracle Turing machine \mathcal{A} will be called a *total algorithm over T* if, for all positive integers ℓ , every model M of T and every $a_1, \dots, a_\ell \in M$, the machine with every input and oracle $\mathcal{O}_M(a_1, \dots, a_\ell)$ is guaranteed to terminate.

4.2. Auxiliary bound and result.

Lemma 4.3. *There is an algorithm that takes as input:*

- language \mathcal{L} ;
- complete recursive theory T given by a Turing machine producing its axioms;
- a total algorithm \mathcal{A} over T ;
- positive integers ℓ and N ;
- a string \mathcal{S} in the input alphabet of \mathcal{A} ;

and computes

- a first-order formula $\varphi = \varphi_{T, \mathcal{A}}(\ell, \mathcal{S}, N)$ in \mathcal{L} in ℓ variables and
- a number $\mathcal{N} := \mathcal{N}_{T, \mathcal{A}}(\ell, \mathcal{S}, N)$

such that, for any model M of T and tuple $\mathbf{a} \in M^\ell$, the following are equivalent:

- (1) the sentence $\varphi(\mathbf{a})$ is true in M ;
- (2) algorithm \mathcal{A} with input \mathcal{S} and oracle $\mathcal{O}_M(\mathbf{a})$ terminates after performing at most N queries to the oracle

and if these statements are true, then the bitsize of the output of \mathcal{A} with input \mathcal{S} and oracle $\mathcal{O}_M(\mathbf{a})$ does not exceed \mathcal{N} .

Proof. We describe an algorithm for computing $\varphi_{T, \mathcal{A}}(\ell, \mathcal{S}, N)$ and $\mathcal{N}_{T, \mathcal{A}}(\ell, \mathcal{S}, N)$. Fix some $\mathcal{L}, T, \mathcal{A}, \ell$, and \mathcal{S} .

We will describe an algorithm that, for a given positive integer s , computes first-order formulas ψ_s and q_s in \mathcal{L} in the variables $\mathbf{x} = (x_1, \dots, x_\ell)$ and a positive integer \mathcal{N}_s such that, for every model M of T and every $\mathbf{a} \in T^\ell$

- $\psi_s(\mathbf{a})$ is true in M iff algorithm \mathcal{A} with input \mathcal{S} and oracle $\mathcal{O}_M(\mathbf{a})$ will perform at least s queries;
- if $\psi_s(\mathbf{a})$ is true in M , then the result of the s -th query will be $q_s(\mathbf{a})$;
- if algorithm \mathcal{A} with input \mathcal{S} and oracle $\mathcal{O}_M(\mathbf{a})$ performs at most s queries, then the bitsize of the output does not exceed \mathcal{N}_s .

Fix some $s \geq 1$ and assume that the algorithm have computed $\psi_1, \dots, \psi_{s-1}$, q_1, \dots, q_{s-1} , and $\mathcal{N}_0, \dots, \mathcal{N}_{s-2}$. Assume that \mathcal{A} with input \mathcal{S} has performed $s-1$ queries. Then whether or not an s -th query will be performed is determined by the results of the first $s-1$ queries. Fix some $\mathbf{r} \in \{\text{True}, \text{False}\}^{s-1}$. It will represent possible results of the first $s-1$ queries. Consider the following formula in \mathcal{L} :

$$\psi_{\mathbf{r}}(\mathbf{x}) := \psi_{s-1}(\mathbf{x}) \wedge \bigwedge_{i=1}^{s-1} (q_i(\mathbf{x}) \iff r_i),$$

where we assume $\psi_0 = \text{True}$. The algorithm uses the recursivity and completeness to check whether the sentence $\exists \mathbf{x} \psi_{\mathbf{r}}(\mathbf{x})$ is false in T [19, Lemma 2.2.8]. If it is, then there is no oracle of the form $\mathcal{O}_M(\mathbf{a})$ such that \mathcal{A} will perform at least $s-1$ queries on it with the results being r_1, \dots, r_{s-1} .

In the case of $\exists \mathbf{x} \psi_{\mathbf{r}}(\mathbf{x})$ is true in T , the algorithm will run \mathcal{A} with input \mathcal{S} and an oracle $\mathcal{O}_{\mathbf{r}}$ that works as follows. For the first $s-1$ queries, $\mathcal{O}_{\mathbf{r}}$ will return r_1, \dots, r_{s-1} . For all subsequent queries, it always returns True. The algorithm will stop the execution of \mathcal{A} if \mathcal{A} makes an s -th query to the oracle, and denote the formula in the query by $q_{\mathbf{r}}$.

Since $\exists \mathbf{x} \psi_{\mathbf{r}}(\mathbf{x})$ is true in T , $\mathcal{O}_{\mathbf{r}}$ gives the same responses to the first $s-1$ queries as some oracle of the form $\mathcal{O}_M(\mathbf{a})$. Since \mathcal{A} must terminate in finite time for every such oracle, one of the following must happen:

- (1) \mathcal{A} will perform an s -th query.
- (2) \mathcal{A} will terminate after performing only $s-1$ queries.

In the former case, as described above, the algorithm will define a formula $q_{\mathbf{r}}$ to be the s -th query. In the latter case, the algorithm will define $\mathcal{N}_{\mathbf{r}}$ to be the bitsize of the output. Then the algorithm computes

$$\begin{aligned} \psi_s(\mathbf{x}) &:= \bigvee_{q_{\mathbf{r}} \text{ is defined}} \psi_{\mathbf{r}}(\mathbf{x}), & q_s(\mathbf{x}) &:= \bigwedge_{q_{\mathbf{r}} \text{ is defined}} (\psi_{\mathbf{r}}(\mathbf{x}) \implies q_{\mathbf{r}}(\mathbf{x})), \\ \mathcal{N}_{s-1} &:= \max \left(\mathcal{N}_{s-2}, \sum_{\mathcal{N}_{\mathbf{r}} \text{ is defined}} \mathcal{N}_{\mathbf{r}} \right), \end{aligned}$$

where we assume $\mathcal{N}_{-1} = -\infty$. If the set $\{\mathbf{r} \mid q_{\mathbf{r}} \text{ is defined}\}$ is empty, the algorithm sets $\psi_s(\mathbf{x}) = \text{False}$ and $q_s(\mathbf{x}) = \text{True}$. Finally, the algorithm returns $\varphi_{T, \mathcal{A}}(\ell, \mathcal{S}, N) := \neg \psi_{N+1}$ and $\mathcal{N}_{T, \mathcal{A}}(\ell, \mathcal{S}, N) := \mathcal{N}_N$. \square

Lemma 4.4. *Let T be a theory and M an \aleph_0 -saturated model. Let $U_1 \supset U_2 \supset U_3 \supset \dots$ be a sequence of definable sets in M^n such that $\bigcap_{i=1}^{\infty} U_i = \emptyset$. Then there exists N such that $U_N = \emptyset$.*

Proof. Assume the contrary, that is, that $U_i \neq \emptyset$ for every $i \geq 1$. We will show that $\bigcap_{i=1}^{\infty} U_i \neq \emptyset$.

We show that a collection of formulas $\{x \in U_i\}_{i=1}^{\infty}$ is finitely satisfiable. Indeed, let $S \subset \mathbb{Z}_{>0}$ be a finite set and $N = \max S$. Then $\bigcap_{i \in S} U_i = U_N \neq \emptyset$. Due to compactness, the countable collection $\{x \in U_i\}_{i=1}^{\infty}$ is satisfiable in some elementary extension of M . Since M is \aleph_0 -saturated, this collection is satisfiable in M . Therefore, $\bigcap_{i=1}^{\infty} U_i \neq \emptyset$. \square

4.3. Main result.

Theorem 4.5. *There exists a computable function $\text{Size}_{T,\mathcal{A}}(\ell, r)$ with input*

- complete recursive theory T ;
- total algorithm \mathcal{A} over T ;
- positive integers ℓ and r

that computes a number N such that for every model M of T , every $\mathbf{a} \in M^\ell$, and every string \mathcal{S} in the alphabet of \mathcal{A} of size at most r the bitsize of the output of \mathcal{A} with input \mathcal{S} and oracle $\mathcal{O}_M(\mathbf{a})$ does not exceed N .

Proof. We will describe an algorithm for computing $\text{Size}_{T,\mathcal{A}}(\ell, r)$. We fix T , \mathcal{A} , ℓ , and r . We will consider \mathcal{S} of length at most r and describe how to compute a bound for the bitsize of the output given that the input is \mathcal{S} . Taking the maximum over all \mathcal{S} of length at most r (there are finitely many of them), we obtain $\text{Size}_{\mathcal{A},T}(\ell, r)$.

The algorithm will compute $\varphi_i := \varphi_{T,\mathcal{A}}(\ell, \mathcal{S}, i)$ for $i = 1, 2, \dots$ using the algorithm from Lemma 4.3. For each φ_i , the algorithm will check whether the formula is equivalent to True in T using the recursivity and completeness [19, Lemma 2.2.8].

If this is true, the algorithm stops and returns $\mathcal{N}_{T,\mathcal{A}}(\ell, \mathcal{S}, i)$ (see Lemma 4.3). It remains to show that the described procedure terminates in finitely many steps. Let M be an \aleph_0 -saturated model of T (it exists, for example, due to [19, Theorem 4.3.12]). For every $i = 1, 2, \dots$, we introduce a definable set

$$U_i := \{\mathbf{a} \in M^\ell \mid \varphi_i(\mathbf{a}) = \text{False}\}.$$

Notice that $U_i = \emptyset$ if and only if $(\varphi_i \iff \text{True})$ in T . Then the definition of φ_i 's implies that $U_1 \supset U_2 \supset \dots$. Assume that $\bigcap_{i=1}^{\infty} U_i$ is not empty and choose an element \mathbf{a} in it. Then \mathcal{A} will not terminate in finitely many steps with input \mathcal{S} and oracle $\mathcal{O}_M(\mathbf{a})$. Thus, $\bigcap_{i=1}^{\infty} U_i = \emptyset$. Lemma 4.4 implies that there exists N such that $U_N = \emptyset$. Then our algorithm will terminate after considering φ_N . \square

5. APPLICATIONS TO DIFFERENTIAL ALGEBRA

In this section, we will apply the results of Section 4 to the theory of differentially closed fields with several commuting derivations.

5.1. Preparation.

Notation 5.1. Let m be a positive integer.

- The language of partial differential rings with m commuting derivations is denoted by $\mathcal{L}_m := \{+, -, \cdot, 0, 1, \partial_1, \dots, \partial_m\}$. We add a separate functional symbol for subtraction for convenience.
- The theory of partial differentially closed fields with m commuting derivations of characteristic zero is denoted by DCF_m . Recall that DCF_m is complete [21, Corollary 3.18] and recursive [21, Section 3.1] (see also [15]).

Notation 5.2. Let m, n, h be positive integers and k a differential field with a set of m commuting derivations $\Delta = \{\partial_1, \dots, \partial_m\}$.

- $\text{Pol}_k(m, n, h)$ denotes the space of all differential polynomials over k in n variables of order at most h and degree at most h .
- The dimension of $\text{Pol}_k(m, n, h)$ (which does not depend on k) will be denoted by $\text{PolDim}(m, n, h)$.

Notation 5.3. Let m, ℓ and n be positive integers.

- Let $\mathcal{L}_m(x_1, \dots, x_\ell)\{y_1, \dots, y_n\}_\Delta$ denote the ring of differential polynomials in differential variables y_1, \dots, y_n with respect to m derivations with the coefficients being terms in the language \mathcal{L}_m in x_1, \dots, x_ℓ (that is, elements of $\mathbb{Z}\{x_1, \dots, x_\ell\}_\Delta$).
- This is a computable differential ring with m commuting derivations. In what follows, we will assume that the algorithms use dense representation to store these polynomials (that is, store all the coefficients up to certain order and certain degree).
- Let k be a differential field with m derivations and $\mathbf{a} \in k^\ell$. Then, for $T \in \mathcal{L}_m(x_1, \dots, x_\ell)\{y_1, \dots, y_n\}_\Delta$, we define $T(\mathbf{a}) \in k\{y_1, \dots, y_n\}_\Delta$ to be the result of evaluating the coefficients of T at \mathbf{a} .

Definition 5.4. A *differential ranking* for $k\{z_1, \dots, z_n\}_\Delta$ is a total order $>$ on $Z := \{\theta z_i \mid \theta \in \Theta_\Delta, 1 \leq i \leq n\}$ satisfying, for all $i, 1 \leq i \leq m$:

- for all $x \in Z$, $\partial_i(x) > x$ and
- for all $x, y \in Z$, if $x > y$, then $\partial_i(x) > \partial_i(y)$.

Notation 5.5. For a Δ -field k and $f \in k\{z_1, \dots, z_n\}_\Delta \setminus k$ and differential ranking $>$,

- $\text{lead}(f)$ is the element of Z of the highest rank appearing in f .
- The leading coefficient of f considered as a polynomial in $\text{lead}(f)$ is denoted by $\text{in}(f)$ and called the initial of f .
- The separant of f is $\frac{\partial f}{\partial \text{lead}(f)}$.
- The rank of f is $\text{rank}(f) = \text{lead}(f)^{\text{deg}_{\text{lead}(f)} f}$. The ranks are compared first with respect to lead , and in the case of equality with respect to deg .
- For $S \subset k\{z_1, \dots, z_n\}_\Delta \setminus k$, the set of initials and separants of S is denoted by H_S .

Definition 5.6 (Characteristic sets).

- For $f, g \in k\{z_1, \dots, z_n\}_\Delta \setminus k$, f is said to be reduced w.r.t. g if no proper derivative of $\text{lead}(g)$ appears in f and $\text{deg}_{\text{lead}(g)} f < \text{deg}_{\text{lead}(g)} g$.
- A subset $\mathcal{A} \subset k\{z_1, \dots, z_n\}_\Delta \setminus k$ is called *autoreduced* if, for all $p \in \mathcal{A}$, p is reduced w.r.t. every element of $\mathcal{A} \setminus \{p\}$. One can show that every autoreduced set is finite [13, Section I.9].
- Let $\mathcal{A} = A_1 < \dots < A_r$ and $\mathcal{B} = B_1 < \dots < B_s$ be autoreduced sets ordered by their ranks (see Notation 5.5). We say that $\mathcal{A} < \mathcal{B}$ if
 - $r > s$ and $\text{rank}(A_i) = \text{rank}(B_i)$, $1 \leq i \leq s$, or
 - there exists q such that $\text{rank}(A_q) < \text{rank}(B_q)$ and, for all $i, 1 \leq i < q$, $\text{rank}(A_i) = \text{rank}(B_i)$.
- An autoreduced subset of the smallest rank of a differential ideal $I \subset k\{z_1, \dots, z_n\}_\Delta$ is called a *characteristic set* of I . One can show that every non-zero differential ideal in $k\{z_1, \dots, z_n\}_\Delta$ has a characteristic set.

- A radical differential ideal I of $k\{z_1, \dots, z_n\}_\Delta$ is said to be *characterizable* if I has a characteristic set C such that $I = \langle C \rangle^{(\infty)} : H_C^\infty$.

Proposition 5.7. *There is a computable function that, for a given positive integer m , computes a total algorithm \mathcal{RG}_m over DCF_m such that, for every differential field k with m derivations and $\mathbf{a} \in k^\ell$, the input-output specification of \mathcal{RG}_m with oracle $\mathcal{O}_k(\mathbf{a})$ is the following:*

Input: *finite subsets A and S of $\mathcal{L}_m(x_1, \dots, x_\ell)\{y_1, \dots, y_n\}_\Delta$;*

Output: *a list of tuples C_1, \dots, C_N from $\mathcal{L}_m(x_1, \dots, x_\ell)\{y_1, \dots, y_n\}_\Delta$ such that*

$$C_1(\mathbf{a}), \dots, C_N(\mathbf{a})$$

is the output of the Rosenfeld-Gröbner algorithm [2, Theorem 9] with input $(A(\mathbf{a}), S(\mathbf{a}))$.

Proof. [2, Theorem 9] states that the only operations performed by the Rosenfeld-Gröbner algorithm with the elements of the ground differential field are arithmetic operations, differentiation, and zero testing. Algorithm \mathcal{RG}_m is constructed to work exactly in the same way as the Rosenfeld-Gröbner algorithm with the only difference that the elements of the ground differential field will be represented as $L(\mathbf{a})$, where $L \in \mathcal{L}_m(x_1, \dots, x_\ell)\{y_1, \dots, y_n\}_\Delta$. The arithmetic operations and differentiations can be performed with L , and zero testing can be performed using the oracle, so \mathcal{RG} will be able to perform the same computations as the Rosenfeld-Gröbner algorithm.

Due to [2, Theorem 5], the Rosenfeld-Gröbner algorithm is guaranteed to terminate on every input. Hence, the same is true for \mathcal{RG}_m . \square

5.2. Bounds.

Theorem 5.8 (Upper bound for Rosenfeld-Gröbner algorithm). *There exists a computable function $\text{RG}(m, n, \ell)$ such that, for every differential field k with m derivations and subsets $A, S \subset \text{Pol}_k(m, n, n)$ with $|A|, |S| \leq \ell$, and every differential ranking, the output of the Rosenfeld-Gröbner algorithm [2, Theorem 9] on A and S will produce at most $\text{RG}(m, n, \ell)$ components with all the orders and degrees not exceeding $\text{RG}(m, n, \ell)$.*

Proof. We fix m , n , and ℓ and compute the total algorithm \mathcal{RG}_m over DCF_m from Proposition 5.7. Let \mathbf{a} be the set of all the coefficients of A and S . Then $|\mathbf{a}| \leq N := 2\ell \text{PolDim}(m, n, n)$. The sets A and S can be presented as evaluations of subsets $\tilde{A}, \tilde{S} \subset \mathcal{L}_m(x_1, \dots, x_N)\{y_1, \dots, y_n\}_\Delta$ at \mathbf{a} such that the orders and degrees of \tilde{A}, \tilde{S} in y_1, \dots, y_n do not exceed n and every coefficient is a single variable x_i . Then the size of (\tilde{A}, \tilde{S}) is bounded by a computable function $S(m, n, N)$.

We run \mathcal{RG}_m with the input $\mathcal{I} = (\tilde{A}, \tilde{S})$ and oracle $\mathcal{O}(\mathbf{a})$. Lemma 4.5 implies that the bitsize of the output will not exceed $\text{Size}_{\mathcal{RG}_m, \text{DCF}_m}(N, S(m, N, n))$.

Since each component takes at least one bit, polynomial of degree d or order d has at least d coefficients (due to the dense representation of the polynomials, see Notation 5.2) requiring at least one bit each, the number of components, the degrees and orders do not exceed the bitsize of the output. Therefore, we can set $\text{RG}(m, n, \ell) = \text{Size}_{\mathcal{RG}, \text{DCF}_m}(N, S(m, N, n))$. \square

Corollary 5.9. *There exists a computable function $\text{CharSet}(m, n, \ell)$ such that, for every computable differential field k with m derivations and subsets $A, S \subset \text{Pol}_k(m, n, n)$ with $|A|, |S| \leq \ell$, and every differential ranking, the ideal*

$\sqrt{\langle A \rangle^{(\infty)} : S^\infty}$ can be written as an intersection of at most $\text{CharSet}(m, n, \ell)$ characterizable differential ideals defined by their characteristic sets with respect to the ranking of order and degree not exceeding $\text{CharSet}(m, n, \ell)$.

Proof. Theorem 5.8 implies that there exists a representation

$$\sqrt{\langle A \rangle^{(\infty)} : S^\infty} = (\langle C_1 \rangle^{(\infty)} : H_{C_1}) \cap \dots \cap (\langle C_N \rangle^{(\infty)} : H_{C_N}),$$

where H_{C_i} is the product of the initials and separants of C_i , and C_i is the characteristic presentation [2, Definition 8] of $\langle C_i \rangle^{(\infty)} : H_{C_i}^\infty$ for every $1 \leq i \leq N$. As noted in [2, p. 108] a characteristic set of $\langle C_i \rangle^{(\infty)} : H_{C_i}^\infty$ can be obtained from C_i by performing reductions until it will become autoreduced. Since differential reduction is a part of the Rosenfeld-Göbner algorithm, it can also be performed by a total algorithm over DCF_m . Therefore, as in the proof of Theorem 5.8, Lemma 4.5 implies that $\langle C_i \rangle^{(\infty)} : H_{C_i}^\infty$ has a characteristic set with degrees and order bounded by a computable function of the degrees and orders of C_i . The latter are bounded by a computable function RG due to Theorem 5.8. Composing these two bounds, we obtain a desired function $\text{CharSet}(m, n, \ell)$. \square

Lemma 5.10. *There exists a computable function $\text{PrimeComp}(m, n)$ such that for every partial differential field k with m derivations, every ranking, and every characterizable differential ideal I defined by a characteristic set $C \subset \text{Pol}_k(m, n, n)$ with respect to this ranking, we have*

- (1) *the number of prime components of I does not exceed $\text{PrimeComp}(m, n)$;*
- (2) *every prime component of I has a characteristic set with respect to the ranking with orders and degrees bounded by $\text{PrimeComp}(m, n)$.*

Proof. Let H be the product of the initials and separants of C . [2, Theorem 4] implies that the number of prime components of $\langle C \rangle^{(\infty)} : H^\infty$ is equal to the number of prime components of the algebraic ideal $(\langle C \rangle^{(\infty)} : H^\infty) \cap R_n$, where R_n is the ring of differential polynomials of order at most n . Since the degrees of elements of C are bounded by n , the Bézout inequality implies that there is a computable bound D for the degree of $I \cap R_n$ in terms of m and n , so this gives a bound for the number of components.

Let P_1, \dots, P_ℓ be the prime components of I . For every $1 \leq i \leq \ell$, $P_i \cap R_n$ is a prime algebraic ideal, and its zero set can be defined by equations of degree at most $\deg(P_i \cap R_n)$ due to [12, Proposition 3]. Therefore, for each $2 \leq i \leq \ell$, we can choose a polynomial in $(P_1 \setminus P_i) \cap R_n$ of degree at most $\deg(P_i \cap R_n)$. Their product Q has degree at most $\deg(I \cap R_n) \leq D$. Observe that

$$P_1 = P_1 : Q^\infty \subset I : Q^\infty = (P_1 : Q^\infty) \cap \dots \cap (P_\ell : Q^\infty) = P_1.$$

Thus, applying Corollary 5.9 to a pair (C, HQ) and using that $|C| \leq \text{PolDim}(m, n)$, we show that P_1 has a characteristic set with orders and degrees bounded by $\text{CharSet}(m, D + n, \text{PolDim}(m, n))$. \square

Theorem 5.11 (Upper bound for the components of a differential variety and their number). *There exists a computable function $\text{Comp}(m, n)$ such that, for all non-negative integers m, n and h and a partial differential field k with m derivations and finite set $F \subset \text{Pol}_k(m, n, h)$:*

- (1) *the number of components in the variety defined by $F = 0$ does not exceed $\text{Comp}(m, \max\{n, h\})$;*

- (2) for every differential ranking and every component X of the variety $F = 0$, X has a characteristic set with respect to the ranking with orders and degrees bounded by $\text{Comp}(m, \max\{n, h\})$.

Proof. Consider any differential ranking. By replacing F with the basis of its linear span, we will further assume that $|F| \leq \text{PolDim}(m, n, h)$ (see Notation 5.2). Corollary 5.9 implies that $\sqrt{\langle F \rangle^{(\infty)}}$ can be represented as an intersection of at most N characterizable ideals with characteristic sets C_1, \dots, C_N of order and degree at most N , where

$$N := \text{CharSet}(m, \max\{n, h\}, \text{PolDim}(m, n, h)).$$

Lemma 5.10 applied to each of C_1, \dots, C_N implies that the number of components of the variety defined by $F = 0$ does not exceed $N \cdot \text{PrimeComp}(m, N)$, and each of them has a characteristic set with orders and degrees not exceeding $\text{PrimeComp}(m, N)$. \square

Remark. It was shown in [11, Theorem 6.1] that there exists a (not necessarily computable) bound for the degrees and orders a characteristic set of a prime differential ideal. The second part of Theorem 5.11 implies that there is a computable bound.

6. APPLICATION TO DELAY PDES

In this section, we will show how Theorem 5.11 applies to the problem of elimination of unknowns in delay PDEs.

6.1. Bounds for Kolchin polynomials for algebraic PDEs.

Definition 6.1. We will say that a Δ -variety $X \subset \mathbb{A}^n$ is *bounded by N* if $N \geq \max(n, m)$ ($m = |\Delta|$) and X can be defined by equations of order and degree at most N .

Notation 6.2. For a numeric polynomial $\omega(t) = \sum_{i=0}^m a_i \binom{t+i}{i}$, we set

$$|\omega| := \sum_{i=0}^m |a_i|.$$

Lemma 6.3. *There exists a computable function $\text{KolchinProj}(N)$ such that for every*

- differential variety $X \subset \mathbb{A}^n$ bounded by N ,
- irreducible component $X_0 \subset X$,
- and linear projection $\pi: \mathbb{A}^n \rightarrow \mathbb{A}^\ell$,

we have $|\omega_Y| \leq \text{KolchinProj}(N)$, where $Y := \overline{\pi(X_0)}^{\text{Kol}}$.

Proof. By performing a linear change of variables, we reduce the problem to the case in which π is the projection to the first ℓ coordinates. Consider a ranking such that

- $x_{\ell+i}$ is greater than every derivative of x_j for every $i > 0$ and $1 \leq j \leq \ell$;
- the restriction of the ranking on x_1, \dots, x_ℓ is an orderly ranking.

Theorem 5.11 implies that X_0 has a characteristic set \mathcal{C} with respect to this ranking with the order bounded by a computable function of N . Since a characteristic set of Y can be obtained from \mathcal{C} by selecting the polynomials only in the first ℓ variables, there is a characteristic set of Y with respect to the orderly ranking with the order bounded by a computable function of N . Then [16, Proposition 3.1] and [16, Fact 2.1] imply that $|\omega_Y|$ is bounded by a computable function of N . \square

Proposition 6.4. *There exists an algorithm that, for every computable function $g(n): \mathbb{Z}_{\geq 0} \rightarrow \mathbb{Z}_{\geq 0}$, produces a number Len_g such that, for every sequence of Kolchin polynomials*

$$\omega_0 > \omega_1 > \dots > \omega_\ell$$

such that $|\omega_i| < g(i)$ for every $0 \leq i \leq \ell$, we have $\ell < \text{Len}_g$.

Proof. By replacing $g(n)$ with $n + \max_{0 \leq k \leq n} g(k)$, we can further assume that $g(n)$ is increasing and $g(n) \geq n$. [22, Definition 2.4.9 and Lemma 2.4.12] define a computable order-preserving map c from the set of all Kolchin polynomials \mathcal{K} to $\mathbb{Z}_{\geq 0}^{m+1}$ (considered with respect to the lexicographic ordering). For $v = (v_0, \dots, v_m) \in \mathbb{Z}_{\geq 0}^{m+1}$, we define $|v| = v_0 + \dots + v_m$. For every function $g: \mathbb{Z}_{\geq 0} \rightarrow \mathbb{Z}_{\geq 0}$, we define

$$\tilde{g}(n) := \max_{\omega \in \mathcal{K}, |\omega| \leq g(n)} |c(\omega)|.$$

Note that if $g(n)$ was computable, then $\tilde{g}(n)$ is also computable.

The sequence $\omega_0 > \omega_1 > \dots$ gives rise to a sequence $c(\omega_0) >_{\text{lex}} c(\omega_1) >_{\text{lex}} \dots$ in $\mathbb{Z}_{\geq 0}^{m+1}$ with $|c(\omega_i)| \leq \tilde{g}(i)$ for every i . [20, Main Lemma] implies that there is an algorithm to compute the maximal length of such a sequence, so there is an algorithm to compute a bound on ℓ from g . \square

6.2. Trains of varieties, partial solutions, and their upper bounds.

Lemma 6.5. *For every Δ - σ -field k of characteristic zero, there exists an extension $k \subset K$ of Δ - σ -fields, where K is a differentially closed Δ - σ^* -field.*

Proof. The proof follows [18, Lemma 6.1] mutatis mutandis and replacing the reference to [3, Theorem 3.15] by [14, Corollary 2.4]. \square

Notation 6.6. Within Sections 6.2 and 6.3, we fix a ground Δ - σ field k and a differentially closed Δ - σ^* -field K given by Lemma 6.5 applied to k . All varieties in Sections 6.2 and 6.3 are considered over K .

Definition 6.7 (Partial solutions).

- For Δ - σ -rings \mathcal{R}_1 and \mathcal{R}_2 , a homomorphism $\phi: \mathcal{R}_1 \rightarrow \mathcal{R}_2$ is called a Δ - σ -homomorphism if, for all i , $\phi \partial_i = \partial_i \phi$ and $\phi \sigma = \sigma \phi$.
- Let \mathcal{R} be a Δ - σ -ring containing a Δ - σ -field k . Let $k[\mathbf{y}_\infty]$ be the Δ - σ -polynomial ring over k in $\mathbf{y} = y_1, \dots, y_r$. Given a point $\mathbf{a} = (a_1, \dots, a_r) \in \mathcal{R}^r$, there exists a unique Δ - σ -homomorphism over k ,

$$\phi_{\mathbf{a}}: k[\mathbf{y}_\infty] \rightarrow \mathcal{R} \quad \text{with} \quad \phi_{\mathbf{a}}(y_i) = a_i \text{ and } \phi_{\mathbf{a}}|_k = \text{id}.$$

Given $f \in k[\mathbf{y}_\infty]$, \mathbf{a} is called a solution of f in \mathcal{R} if $f \in \text{Ker}(\phi_{\mathbf{a}})$.

- For a Δ - σ - k -algebra \mathcal{R} and $I = \mathbb{N}$ or \mathbb{Z} , the sequence ring \mathcal{R}^I has the following structure of a Δ - σ -ring (Δ - σ^* -ring for $I = \mathbb{Z}$) with σ and Δ defined by

$$\sigma((x_i)_{i \in I}) := (x_{i+1})_{i \in I} \quad \text{and} \quad \partial_j((x_i)_{i \in I}) := (\partial_j(x_i))_{i \in I}.$$

For a k - Δ - σ -algebra \mathcal{R} , \mathcal{R}^I can be considered a k - Δ - σ -algebra by embedding k into \mathcal{R}^I in the following way:

$$a \mapsto (\sigma^i(a))_{i \in I}, \quad a \in k.$$

For $f \in k[\mathbf{y}_\infty]$, a solution of f with components in \mathcal{R}^I is called a *sequence solution of f in \mathcal{R}* .

- Given $f \in \mathcal{R}[\mathbf{y}_\infty]$, the order of f is defined to be the maximal $\text{ord } \theta + j$ such that $\theta \sigma^j y_k$ effectively appears in f for some k , denoted by $\text{ord}(f)$.
- The relative order of f with respect to Δ (resp. σ), denoted by $\text{ord}_\Delta(f)$ (resp. $\text{ord}_\sigma(f)$), is defined as the maximal $\text{ord } \theta$ (resp. j) such that $\theta \sigma^j y_k$ effectively appears in f for some k .
- Let $F = \{f_1, \dots, f_N\} \subset k[\mathbf{y}_\infty]$, where $\mathbf{y} = y_1, \dots, y_r$, be a set of Δ - σ -polynomials. Suppose $h = \max\{\text{ord}_\sigma(f) \mid f \in F\}$. A sequence of tuples $(\bar{a}_1, \dots, \bar{a}_r) \in K^{\ell+h} \times \dots \times K^{\ell+h}$ is called a *partial solution* of F of length ℓ if $(\bar{a}_1, \dots, \bar{a}_r)$ is a Δ -solution of the system in $\mathbf{y}_{\infty, \ell+h-1}$:

$$\{\sigma^i(F) = 0 \mid 0 \leq i \leq \ell - 1\}.$$

We associate the following geometric data with the above set F of Δ - σ -polynomials:

- the Δ -variety $X \subset \mathbb{A}^H$ defined by $f_1 = 0, \dots, f_N = 0$ regarded as Δ -equations in $k[\mathbf{y}_{\infty, h}]$ with $H = r(h+1)$, and
- two projections $\pi_1, \pi_2 : \mathbb{A}^H \rightarrow \mathbb{A}^{H-r}$ defined by

$$\begin{aligned} \pi_1(a_1, \dots, \sigma^h(a_1); \dots; a_r, \dots, \sigma^h(a_r)) \\ &:= (a_1, \sigma(a_1), \dots, \sigma^{h-1}(a_1); \dots; a_r, \dots, \sigma^{h-1}(a_r)), \\ \pi_2(a_1, \dots, \sigma^h(a_1); \dots; a_r, \dots, \sigma^h(a_r)) \\ &:= (\sigma(a_1), \dots, \sigma^h(a_1); \dots; \sigma(a_r), \dots, \sigma^h(a_r)). \end{aligned}$$

Let $\sigma(X)$ denote the Δ -variety in \mathbb{A}^H defined by $f_1^\sigma, \dots, f_N^\sigma$, where f_i^σ is the result by applying σ to the coefficients of f_i .

Definition 6.8. A sequence $p_1, \dots, p_\ell \in \mathbb{A}^H$ is a *partial solution of the triple* (X, π_1, π_2) if

- (1) for all i , $1 \leq i \leq \ell$, we have $p_i \in \sigma^{i-1}(X)$ and
- (2) for all i , $1 \leq i < \ell$, we have $\pi_1(p_{i+1}) = \pi_2(p_i)$.

A two-sided infinite sequence with such a property is called a *solution of the triple* (X, π_1, π_2) .

Lemma 6.9. *For every positive integer ℓ , F has a partial solution of length ℓ if and only if the triple (X, π_1, π_2) has a partial solution of length ℓ . The system F has a solution in $K^{\mathbb{Z}}$ if and only if the triple (X, π_1, π_2) has a solution.*

Proof. As in [18, Lemma 6.5]. □

Definition 6.10. For $\ell \in \mathbb{N}$ or $+\infty$, a sequence of irreducible Δ -subvarieties (Y_1, \dots, Y_ℓ) in \mathbb{A}^H is said to be a *train of length ℓ in X* if

- (1) for all i , $1 \leq i \leq \ell$, we have $Y_i \subseteq \sigma^{i-1}(X)$ and
- (2) for all i , $1 \leq i < \ell$, we have $\overline{\pi_1(Y_{i+1})}^{\text{Kol}} = \overline{\pi_2(Y_i)}^{\text{Kol}}$.

Lemma 6.11. *For every train (Y_1, \dots, Y_ℓ) in X , there exists a partial solution p_1, \dots, p_ℓ of (X, π_1, π_2) such that for all i , we have $p_i \in Y_i$. In particular, if there is an infinite train in X , then there is a solution of the triple (X, π_1, π_2) .*

Proof. As in [18, Lemma 6.7]. \square

For two trains $Y = (Y_1, \dots, Y_\ell)$ and $Y' = (Y'_1, \dots, Y'_\ell)$, denote $Y \subseteq Y'$ if $Y_i \subseteq Y'_i$ for each i . Given an increasing chain of trains $Y_i = (Y_{i,1}, \dots, Y_{i,\ell})$,

$$\left(\overline{\bigcup_i Y_{i,1}}^{\text{Kol}}, \dots, \overline{\bigcup_i Y_{i,\ell}}^{\text{Kol}} \right)$$

is a train in X that is an upper bound for this chain. (For each j , $\overline{\bigcup_i Y_{i,j}}^{\text{Kol}}$ is an irreducible δ -variety in $\sigma^{j-1}(X)$.) So by Zorn's lemma, maximal trains of length ℓ always exist in X .

For $\ell \in \mathbb{N}$, consider the product

$$\mathbf{X}_\ell := X \times \sigma(X) \times \dots \times \sigma^{\ell-1}(X)$$

and denote the projection of \mathbf{X}_ℓ onto $\sigma^{i-1}(X)$ by $\varphi_{\ell,i}$. Let

$$\mathbf{W}_\ell(X, \pi_1, \pi_2) := \{p \in \mathbf{X}_\ell : \pi_2(\varphi_{\ell,i}(p)) = \pi_1(\varphi_{\ell,i+1}(p)), i = 1, \dots, \ell - 1\}.$$

Lemma 6.12. *Let (X, π_1, π_2) be a triple with X bounded by n . Then, for every ℓ , the number of maximal trains of length ℓ in X does not exceed $\text{Comp}(m, \ell n)$.*

Proof. This follows from Theorem 5.11 and rewritten mutatis mutandis for several commuting derivations [18, Lemma 6.8]. \square

Definition 6.13. Let (X, π_1, π_2) be a triple and $\omega(t)$ be a numeric polynomial. We define $B(X, \omega) \in \mathbb{Z} \cup \{\infty\}$ as the smallest value that is greater than the length of any train in X with Kolchin polynomials at least ω .

Lemma 6.14. *Let X be a differential variety bounded by n such that $B(X, 0) < \infty$. Then $B(X, \omega_X)$ does not exceed the number of components of X plus one.*

Proof. Denote the number of components in X by N and assume that there is a train (Y_1, \dots, Y_{N+1}) with the Kolchin polynomial at least ω_X . Then each of $Y_1, \sigma^{-1}(Y_2), \dots, \sigma^{-N}(Y_{N+1})$ must be a component of X , so there exist $1 \leq i < j \leq N + 1$ such that $Y_j = \sigma^{j-i}Y_i$. Thus, there exists an infinite train $(Y_1, \dots, Y_i, Y_{i+1}, \dots, Y_{j-1}, \sigma^{j-i}(Y_i), \sigma^{j-i}(Y_{i+1}), \dots)$ in X . This contradicts to $B(X, 0) < \infty$. \square

Lemma 6.15. *There exists a computable function $\text{Iter}(n, D)$ such that, for every triple (X, π_1, π_2) such that*

- $B(X, 0) < \infty$
- X is bounded by n

and every numeric polynomial $\omega_1(t) > 0$, there exists a numeric polynomial $\omega_2(t) \geq 0$ such that

- $\omega_2(t) < \omega_1(t)$;
- $|\omega_2| \leq \text{Iter}(n, B(X, \omega_1))$;
- $B(X, \omega_2) \leq \text{Iter}(n, B(X, \omega_1))$.

Proof. The proof follows [18, Lemma 6.20]. Let $B_1 := B(X, \omega_1)$, and let T be the number of maximal trains of length B_1 in X . We set $B_2 := B_1 + T$. Lemma 6.12 implies that T is bounded by $\text{Comp}(m, nB_1)$. Consider the fibered

product $\mathbf{W}_{B_1}(X, \pi_1, \pi_2)$, and, for each irreducible components W in it, denote the corresponding train by Y_W . We set (assuming $\max \emptyset = 0$)

$$\omega_2 := \max\{\omega_{Y_W} \mid \omega_{Y_W} < \omega_1, W \text{ is a component of } \mathbf{W}_{B_1}\}.$$

We will show that $B(X, \omega_2) \leq B_1 + T$. Assume that there is a maximal train (Y_1, \dots, Y_{B_2}) in X with the Kolchin polynomial at least ω_2 . Introduce $T + 1$ trains $Z^{(1)}, \dots, Z^{(T+1)}$ of length B_1 in $X, \sigma(X), \dots, \sigma^T(X)$, respectively, such that for each j ,

$$Z^{(j)} = (Z_1^{(j)}, \dots, Z_T^{(j)}) := (Y_j, \dots, Y_{j+B_1-1}).$$

Then for each j , consider a maximal train $\tilde{Z}^{(j)}$ of length B_1 containing $Z^{(j)}$. So $\sigma^{-j+1}(\tilde{Z}^{(j)})$ is a maximal train of length B_1 in X . There are two cases to consider:

$$\text{(Case 1)} \quad \{\omega_{Y_W}(t) \mid \omega_{Y_W}(t) < \omega_1(t), W \text{ is a component of } \mathbf{W}_{B_1}\} = \emptyset.$$

In this case, $\tilde{Z}^{(1)}$ is a train in X with Kolchin polynomial at least ω_1 . This contradicts the definition of $B(X, \omega_1)$.

$$\text{(Case 2)} \quad \{\omega_{Y_W}(t) \mid \omega_{Y_W}(t) < \omega_1(t), W \text{ is a component of } \mathbf{W}_{B_1}\} \neq \emptyset.$$

By the definition of $B(X, \omega_1)$, for every j , $\omega_{\sigma^{-j+1}(\tilde{Z}^{(j)})}(t) < \omega_1(t)$. This implies that, for each j ,

$$\omega_{\sigma^{-j+1}(\tilde{Z}^{(j)})}(t) = \omega_2(t).$$

Since there are only T maximal trains in X of length B_1 , there exist $a < b$ such that

$$\sigma^{-a+1}(\tilde{Z}^{(a)}) = \sigma^{-b+1}(\tilde{Z}^{(b)}) =: Z.$$

Since $\omega_Z = \omega_2$, there exists ℓ such that $\omega_{Z_\ell} = \omega_2$. Since

$$\omega_{\sigma^{-a+1}(Z_\ell^{(a)})} = \omega_2 \quad \text{and} \quad \sigma^{-a+1}(Z_\ell^{(a)}) \subseteq Z_\ell$$

we have $\sigma^{-a+1}(Z_\ell^{(a)}) = Z_\ell$. Similarly, we can show $\sigma^{-b+1}(Z_\ell^{(b)}) = Z_\ell$. Hence,

$$\sigma^{-a+1}(Y_{a+\ell-1}) = \sigma^{-a+1}(Z_\ell^{(a)}) = \sigma^{-b+1}(Z_\ell^{(b)}) = \sigma^{-b+1}(Y_{b+\ell-1}).$$

Thus, we have $Y_{b+\ell-1} = \sigma^{b-a}(Y_{a+\ell-1})$. This contradicts the fact that $B(X, 0) < \infty$.

It remains to show that $|\omega_2|$ is bounded by a computable function of n and B_1 . Let W be a component of \mathbf{W}_{B_1} such that $\omega_{Y_W} = \omega_2$. Let $Y_W = (Y_{W,1}, \dots, Y_{W,B_1})$. There exists $1 \leq i \leq B_1$ such that $\omega_{Y_i} = \omega_2$. Since Y_i is the Kolchin closure of a linear projection of a component of \mathbf{W}_{B_1} and \mathbf{W}_{B_1} is bounded by $B_1 n$, Lemma 6.3 implies that $|\omega_2|$ is bounded by a computable function of n and B_1 .

Taking $\text{Iter}(n, D)$ to be the maximum of the computable bounds for $B(X, \omega_2)$ and $|\omega_2|$, we conclude the proof. \square

Definition 6.16. Let n be a positive integer and $\omega(t)$ be a numeric polynomial such that $\omega > 0$. We define $B(n, \omega) \in \mathbb{Z} \cup \{\infty\}$ as the smallest value such that, for every affine differential variety X bounded by n , if there exists a train in X with Kolchin polynomial at least ω of length at least $B(n, \omega)$, then there exists an infinite train in X .

Proposition 6.17. $B(n, 0)$ is bounded by a computable function $A(n)$.

Proof. We recursively define the following function $G(n)$ on nonnegative integers

$$\begin{aligned} G(0) &:= \max(\text{Components}(n) + 1, \text{KolchinProj}(n)), \\ G(j+1) &:= \text{Iter}(n, G(j)), \quad j \geq 0. \end{aligned}$$

Consider a variety X bounded by n such that there is no infinite train in X , that is $B(X, 0) < \infty$. Lemma 6.14 implies that $B(X, \omega_X) - 1$ does not exceed the number of components of X . Hence Theorem 5.11 implies that $B(X, \omega_X) \leq \text{Comp}(n) + 1$. Lemma 6.3 implies that $|\omega_X| \leq \text{KolchinProj}(n)$. Repeatedly applying Lemma 6.15, we obtain a sequence of numeric polynomials

$$\omega_0 := \omega_X > \omega_1 > \omega_2 > \dots$$

such that, for every $1 \leq i \leq L$, we have $B(X, \omega_i) \leq G(i)$ and $|\omega_i| \leq G(i)$. Since the Kolchin polynomials are well-ordered, there exists L such that $\omega_L = 0$. Proposition 6.4 implies that $L \leq \text{Len}_G$. Hence, $B(X, 0) \leq G(\text{Len}_G)$, where the right-hand side is a computable function of n . Set $A(n) := G(\text{Len}_G)$, then $B(n, 0) \leq A(n)$. \square

Corollary 6.18. *For all r, m and $s \in \mathbb{Z}_{\geq 0}$, and a set of Δ - σ polynomials $F \subset k[\mathbf{y}_s]$ with $|\Delta| = m$, $\deg F \leq s$ and $|\mathbf{y}| = r$, $F = 0$ has a solution in $K^{\mathbb{Z}}$ if and only if $F = 0$ has a partial solution of computable length $A(\max\{r, m, s\})$.*

Proof. As in [18, Corollary 6.21]. \square

6.3. Upper bound for delay PDEs.

Theorem 6.19. *For all non-negative integers r, m , and s , there exists a computable $B = B(r, m, s)$ such that, for all:*

- non-negative integers q and t ,
- Δ - σ -fields k with $\text{char } k = 0$ and $|\Delta| = m$,
- sets of Δ - σ -polynomials $F \subset k[\mathbf{x}_t, \mathbf{y}_s]$, where $\mathbf{x} = x_1, \dots, x_q$, $\mathbf{y} = y_1, \dots, y_r$, and $\deg_{\mathbf{y}} F \leq s$,

we have

$$\langle \sigma^i(F) \mid i \in \mathbb{Z}_{\geq 0} \rangle^{(\infty)} \cap k[\mathbf{x}_{\infty}] \neq \{0\} \iff \langle \sigma^i(F) \mid i \in [0, B] \rangle^{(B)} \cap k[\mathbf{x}_{B+t}] \neq \{0\}.$$

Proof. The proof closely follows [18, Theorem 6.22]. The “ \Leftarrow ” implication is straightforward. We will prove the “ \Rightarrow ” implication. For this, let $A := A(\max\{r, m, s\})$ from Corollary 6.18, and let B be a computable bound obtained from [10, Theorem 3.4] with

$$m \leftarrow m, \quad n \leftarrow r(A + s + 1), \quad h \leftarrow s, \quad \text{and} \quad D \leftarrow s.$$

By assumption,

$$(1) \quad 1 \in \langle \sigma^i(F) \mid i \in \mathbb{Z}_{\geq 0} \rangle^{(\infty)} \cdot k(\mathbf{x}_{\infty})[\mathbf{y}_{\infty}].$$

Suppose that

$$(2) \quad \langle \sigma^i(F) \mid i \in [0, A] \rangle^{(B)} \cap k[\mathbf{x}_{B+t}] = \{0\}.$$

If

$$1 \in \langle \sigma^i(F) \mid i \in [0, A] \rangle^{(B)} \cdot k(\mathbf{x}_{B+t})[\mathbf{y}_{\infty, A+s}],$$

then there would exist $c_{i,j} \in k(\mathbf{x}_{B+t})[\mathbf{y}_{\infty, A+s}]$ such that

$$(3) \quad 1 = \sum_{\theta \in \Theta_{\Delta}(B)} \sum_{j=0}^A \sum_{f \in F} c_{i,j} \theta(\sigma^j(f)).$$

Multiplying equation (3) by the common denominator in the variables \mathbf{x}_{B+t} , we obtain a contradiction with (2). Hence, by [10, Theorem 3.4],

$$1 \notin \langle \sigma^i(F) \mid i \in [0, A] \rangle^{(\infty)} \cdot k(\mathbf{x}_{B+t})[\mathbf{y}_{\infty, A+s}].$$

By Lemma 6.5, there exists a differentially closed Δ - σ^* -field extension $L \supset k(\mathbf{x}_{\infty}) \supset k(\mathbf{x}_{B+t})$. Then differential Nullstellensatz implies that the system of differential equations

$$\{\sigma^i(F) = 0 \mid i \in [0, A]\}$$

in the unknowns $\mathbf{y}_{\infty, A+s}$ has a solution in L . Then the system $F = 0$ has a partial solution of length $A + 1$ in L . Now from (1), we see that the system $F = 0$ has no solutions in $L^{\mathbb{Z}}$. Together with the existence of a partial solution of length $A + 1$, this contradicts to Corollary 6.18. \square

ACKNOWLEDGMENTS

This work was partially supported by the NSF grants CCF-1564132, CCF-1563942, DMS-1760448, DMS-1760413, DMS-1853650, DMS-1853482, and DMS-1800492; and the NSFC grants (11971029, 11688101).

REFERENCES

- [1] T. Bächler, V. Gerdt, M. Lange-Hegermann, and D. Robertz. Algorithmic Thomas decomposition of algebraic and differential systems. *Journal of Symbolic Computation*, 47(10):1233–1266, 2012. URL <https://doi.org/10.1016/j.jsc.2011.12.043>.
- [2] F. Boulier, D. Lazard, F. Ollivier, and M. Petitot. Computing representations for radicals of finitely generated differential ideals. *Applicable Algebra in Engineering, Communication and Computing*, 20:73–121, 2009. URL <https://doi.org/10.1007/s00200-009-0091-7>.
- [3] R. F. Bustamante Medina. Differentially closed fields of characteristic zero with a generic automorphism. *Revista de Matematica: Teoria y Aplicaciones*, 14(1):81–100, 2007. URL <https://doi.org/10.15517/rmta.v14i1.282>.
- [4] O. Chapuis and P. Koiran. Saturation and stability in the theory of computation over the reals. *Annals of Pure and Applied Logic*, 99(1):1–49, 1999. URL [https://doi.org/10.1016/S0168-0072\(98\)00060-8](https://doi.org/10.1016/S0168-0072(98)00060-8).
- [5] X.-S. Gao, Y. Luo, and C. Yuan. A characteristic set method for ordinary difference polynomial systems. *Journal of Symbolic Computation*, 44(3):242–260, 2009. URL <https://doi.org/10.1016/j.jsc.2007.05.005>.
- [6] X.-S. Gao, J. Van der Hoeven, C.-M. Yuan, and G.-L. Zhang. Characteristic set method for differential-difference polynomial systems. *Journal of Symbolic Computation*, 44(9):1137–1163, 2009. URL <https://doi.org/10.1016/j.jsc.2008.02.010>.
- [7] V. Gerdt and D. Robertz. Algorithmic approach to strong consistency analysis of finite difference approximations to PDE systems. In *ISSAC '19: Proceedings of the 2019 on International Symposium on Symbolic and Algebraic Computation*, pages 163–170. ACM Press, 2019. URL <https://doi.org/10.1145/3326229.3326255>.
- [8] O. Golubitsky, M. Kondratieva, A. Ovchinnikov, and A. Szanto. A bound for orders in differential Nullstellensatz. *Journal of Algebra*, 322(11):3852–3877, 2009. URL <https://doi.org/10.1016/j.jalgebra.2009.05.032>.

- [9] J. B. Goode. Accessible telephone directories. *The Journal of Symbolic Logic*, 59:92–105, 1994. URL <https://www.jstor.org/stable/2275252>.
- [10] R. Gustavson, M. Kondratieva, and A. Ovchinnikov. New effective differential Nullstellensatz. *Advances in Mathematics*, 290:1138–1158, 2016. URL <http://dx.doi.org/10.1016/j.aim.2015.12.021>.
- [11] M. Harrison-Trainor, J. Klys, and R. Moosa. Nonstandard methods for bounds in differential polynomial rings. *Journal of Algebra*, 360:71–86, 2012. URL <https://doi.org/10.1016/j.jalgebra.2012.03.013>.
- [12] J. Heintz. Definability and fast quantifier elimination in algebraically closed fields. *Theoretical Computer Science*, 24(3):239–277, 1983. URL [http://dx.doi.org/10.1016/0304-3975\(83\)90002-6](http://dx.doi.org/10.1016/0304-3975(83)90002-6).
- [13] E. Kolchin. *Differential Algebra and Algebraic Groups*. Academic Press, New York, 1973.
- [14] O. León Sánchez. On the model companion of partial differential fields with an automorphism. *Israel Journal of Mathematics*, 212:419–442, 2016. URL <https://doi.org/10.1007/s11856-016-1292-y>.
- [15] O. León Sánchez. Algebro-geometric axioms for $\text{DCF}_{m,0}$. *Fundamenta Mathematicae*, 243:1–8, 2018. URL <https://doi.org/10.4064/fm228-11-2017>.
- [16] O. León Sánchez. Estimates for the coefficients of differential dimension polynomials. *Mathematics of Computation*, 88:2959–2985, 2019. URL <https://doi.org/10.1090/mcom/3429>.
- [17] O. León Sánchez and R. Moosa. The model companion of differential fields with free operators. *Journal of Symbolic Logic*, 81(2):493–509, 2016. URL <https://doi.org/10.1017/jsl.2015.76>.
- [18] W. Li, A. Ovchinnikov, G. Pogudin, and T. Scanlon. Elimination of unknowns for systems of algebraic differential-difference equations, 2018. URL <https://arxiv.org/abs/1812.11390>.
- [19] D. Marker. *Model theory: An introduction*. Springer-Verlag New York, 2002. URL <https://doi.org/10.1007/b98860>.
- [20] K. McAloon. Petri nets and large finite sets. *Theoretical Computer Science*, 32(1-2):173–183, 1984. URL [https://doi.org/10.1016/0304-3975\(84\)90029-X](https://doi.org/10.1016/0304-3975(84)90029-X).
- [21] T. McGrail. The model theory of differential fields with finitely many commuting derivations. *Journal of Symbolic Logic*, 65(2):885–913, 2000. URL <https://doi.org/10.2307/2586576>.
- [22] A. V. Mikhalev, A. Levin, E. Pankratiev, and M. Kondratieva. *Differential and Difference Dimension Polynomials*, volume 461 of *Mathematics and Its Applications*. Springer Netherlands, 1999. URL <https://doi.org/10.1007/978-94-017-1257-6>.
- [23] R. Moosa and T. Scanlon. Model theory of fields with free operators in characteristic zero. *Journal of Mathematical Logic*, 14(2):1450009, 2014. URL <https://doi.org/10.1142/s0219061314500093>.
- [24] C. H. Papadimitriou. *Computational Complexity*. Pearson, 1993.
- [25] W. Simmons and H. Towsner. Proof mining and effective bounds in differential polynomial rings. *Advances in Mathematics*, 343:567–623, 2019. URL <https://doi.org/10.1016/j.aim.2018.11.026>.
- [26] L. van den Dries and K. Schmidt. Bounds in the theory of polynomial rings over fields. a nonstandard approach. *Inventiones mathematicae*, 76:77–91, 1984.

URL <https://doi.org/10.1007/BF01388493>.

KLMM, ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES,
NO.55 ZHONGGUANCUN EAST ROAD, 100190, BEIJING, CHINA

E-mail address: liwei@mmrc.iss.ac.cn

CUNY QUEENS COLLEGE, DEPARTMENT OF MATHEMATICS, 65-30 KISSENA BLVD, QUEENS, NY
11367, USA AND CUNY GRADUATE CENTER, MATHEMATICS AND COMPUTER SCIENCE, 365 FIFTH
AVENUE, NEW YORK, NY 10016, USA

E-mail address: aovchinnikov@qc.cuny.edu

ÉCOLE POLYTECHNIQUE, INSTITUT POLYTECHNIQUE DE PARIS, LIX, CNRS, 1 RUE HONORÉ
D'ESTIENNE D'ORVES, 91120, PALAISEAU, FRANCE AND NATIONAL RESEARCH UNIVERSITY HIGHER
SCHOOL OF ECONOMICS, MOSCOW, RUSSIA

E-mail address: gleb.pogudin@polytechnique.edu

UNIVERSITY OF CALIFORNIA, BERKELEY, DEPARTMENT OF MATHEMATICS, BERKELEY, CA
94720-3840, USA

E-mail address: scanlon@math.berkeley.edu